



United Nations
Environment
Programme



Distr.
LIMITED

UNEP/WG.7/14
25 February 1977

Original: ENGLISH

UNEP Meeting of Experts,
designated by governments, intergovernmental
and non-governmental organizations,
on the Ozone Layer

Washington, D.C., 1 - 9 March 1977

THE EFFECT OF CHLOROFLUOROCARBONS ON THE

CONCENTRATION OF ATMOSPHERIC OZONE

submitted by

THE INTERNATIONAL CHAMBER OF COMMERCE

on behalf of

MANUFACTURING CHEMISTS ASSOCIATION

Review of Industry-Sponsored Research

THE EFFECT OF CHLOROFLUOROCARBONS ON THE CONCENTRATION OF ATMOSPHERIC OZONE

Prepared by: J. R. Soulen

December 31, 1976

This review covers the research program funded by the six U. S. and 14 foreign manufacturers of chlorofluorocarbons that is administered by the Manufacturing Chemists Association and the research undertaken internally by several of the companies.

Although there has been little new scientific information that would change the conclusions of the National Academy of Sciences reports^{1,2} which were issued in September, there is continued high interest in the chlorofluorocarbon-ozone issue on the part of government agencies, industry, and certain public interest groups. It is therefore worthwhile to review the present gaps in our scientific knowledge, and the steps that are being taken to reduce them in industry-sponsored research programs. This is best done using the framework of the NAS Panel report.¹ It will be recalled that the prediction of 2-20% steady state ozone depletion, which would not be achieved for ca. 100 years, was calculated from known data and particular assumptions for (1) release rates of FC-11 and FC-12, (2) atmospheric transport of these molecules, and (3) stratospheric chemistry. The existence of as-yet-unknown factors that could substantially alter this ozone depletion estimate was clearly recognized by the NAS Panel. Indeed, their first recommendation

1. "Halocarbons: Effects on Stratospheric Ozone," Panel on Atmospheric Chemistry, National Academy of Sciences, September 13, 1976.

2. "Halocarbons: Environmental Effects of Chlorofluoromethane Release," Committee on Impacts of Stratospheric Change, National Academy of Sciences, September 13, 1976.

for future work reads:

"In general terms, the greatest need is for verification of stratospheric ozone chemistry thru measurements of trace atmospheric constituents, both stable and reactive. The dearth of such measurements is a serious limitation to establishing the reliability of calculations and predictions of stratospheric effects."³

Dr. Tukey, in his statement before the Subcommittee on the Upper Atmosphere of the U. S. Senate December 15, reemphasized the two kinds of uncertainties in the NAS reports: (1) uncertainties associated with imprecise knowledge of numerical data, which necessitated giving the predicted ozone depletion as a range of possible values from 2-20%, and (2) presently unknown factors that could drastically alter the predicted depletion, possibly taking it below the 2-20% range. As stated in the NAS Committee report, "This range does not allow for possible inadequacies in the bases of the calculation," including "essential chemical reactions not so far recognized as such" and "the possibility of unexpected effects of tropospheric sinks."⁴

Stratospheric chemistry related to the chlorofluorocarbon-ozone question was felt to be particularly uncertain by the NAS Panel. Earlier this year the molecule ClONO_2 , which was dismissed in the earliest considerations of the fluorocarbon-ozone question, was again brought up. The NAS Panel determined that inclusion of ClONO_2 chemistry would reduce ozone depletion estimates by a factor of 1.85. ClONO_2 is just one example of how new chemistry could affect the calculated amount of ozone depletion strongly. As stated in the NAS Committee report, "The probable importance of ClONO_2 was a surprise. Whether or not there is another such surprise in store remains to be seen."⁵

3. Reference 1, page F-1a.

4. Reference 2, pages 1-3,4.

5. Reference 2, page 5-12.

The NAS Panel listed six specific objectives to reduce the uncertainty in ozone depletion estimates due both to factors already known and to the possibility of finding subsequently an important unidentified factor. The most important studies, expected to require one to two years, are as follows:

1. "Establish the role of ClONO₂ in the stratosphere--This might be accomplished by comparing infrared measurements of its concentration with calculated results. However, particularly if disparities are revealed by such comparisons, there will need to be further study of the stratospheric processes that might convert the ClONO₂ back into catalytically active species."⁶

The chlorofluorocarbon industry is supporting the work of Dr. D. G. Murcray of the University of Denver in his efforts to determine the stratospheric concentration of ClONO₂. This support is for balloon flights to make measurements by infrared absorption spectroscopy, laboratory studies to obtain the necessary calibration spectra for ClONO₂, and assistance in the preparative work to obtain samples for the laboratory studies. Several more months will be required before definitive information will be available from flights carried out in 1976. On the basis of indications to date, significant amounts of ClONO₂ in the stratosphere are neither proved nor ruled out.

A major problem in positive identification of molecules using the infrared technique and in determining their concentration is possible interferences from other molecules that absorb in the same spectral region. Greater resolution can minimize this problem and the chlorofluorocarbon manufacturers are supporting Dr. H. J. Buijs of Bomem, Inc., to construct a new spectrometer with an anticipated resolution of 0.01 cm⁻¹. This instrument will be useful in the wavelength region of interest for ClONO₂ and should be completed in early 1977.

6. Reference 1, page F-2.

For long term monitoring of stratospheric molecules, continuous ground-based observations will be much more economical than in situ measurements from balloons, airplanes, or rockets. Scientists from the Du Pont Company and the National Bureau of Standards have collaborated to determine the microwave spectrum of ClONO_2 ⁷, which will be useful for both air and ground based measurements by millimeter wavelength techniques.

The rates of formation and destruction of ClONO_2 are important to the understanding of its stratospheric chemistry, as is knowledge of its mechanism of decomposition (e.g., photolysis, reaction) and the products of its decomposition. The chlorofluorocarbon industry is sponsoring research at the University of Illinois under Dr. J. W. Birks, who determined the rate of formation of ClONO_2 via the reaction $\text{ClO} + \text{NO}_2 + \text{M} \rightarrow \text{ClONO}_2 + \text{M}$ (where M is a molecule such as N_2 or He). Dr. Birks is also studying the rate of destruction of ClONO_2 by both homogeneous and heterogeneous reactions that could result in removal of reactive species from the catalytic ozone destruction cycle.

The photochemistry of ClONO_2 that is relevant to its stratospheric behavior can also be studied in the laboratory. The chlorofluorocarbon manufacturers are supporting work by Dr. J. Wiesenfeld at Cornell University and Dr. R. B. Timmons of Catholic University to determine photoabsorption characteristics and photodecomposition products of ClONO_2 as a function of wavelength and temperature.

2. "Verify the reliability of predicted ozone reductions--By this we mean a more direct means than has been available so far for estimating and narrowing the total uncertainty of the ozone reductions predicted for the CFMs, including the possibility of as yet unidentified processes. Observations of ClO and Cl profiles in the stratosphere and their comparison with calculated values seem to be the most likely means of accomplishing this objective. However, the analysis of any disparities probably will be difficult, and requires comparisons with simultaneous measurements of other species such as HCl and O_3 . Preferably,

7. R. D. Suenram, D. R. Johnson, L. C. Glasgow, and P. Z. Meakin, Geophys. Res. Lett. **3**, 611-4, October, 1976.

the observations of C10 and C1 should be by two independent methods, such as resonance fluorescence and microwave emission in the case of C10."⁸

An approach to verify predicted ozone reductions not emphasized in the shorter-term objectives of the NAS Panel is that based on direct observations of stratospheric ozone itself. Dr. William J. Hill and his colleagues at Allied Chemical Corporation have applied the technique of time series analysis to data from nine long-term ozone stations well-spaced around the world. Their conclusions⁹ are of considerable importance:

1. No detectable abnormal decline in ozone has occurred during the six-year period 1970-75.
2. The predicted cumulative ozone depletions due to FC-11 and FC-12, subsonic and supersonic aircraft, bromine compounds, nitrogenous fertilizers, and a number of halocarbons, when treated cumulatively appear to be too large.
3. The calculated upper limit of 20% steady state depletion (ca. 100 years hence) due to FC-11 and FC-12 appears very unlikely.
4. The time series analysis of data from nine globally representative stations could detect a 1.5% abnormal ozone change. Appropriate action following this early warning could limit the maximum reversible change to only 2.3%. Use of data from a larger number of ground based or satellite stations could increase the detectability and decrease the impact on the ozone layer.

The chlorofluorocarbon manufacturers are supporting Drs. E. Parzen and M. Pagano of the State University of New York at Buffalo to carry out further research on time-series analysis of ozone data. Work of this type clearly can have great importance to both the stratospheric ozone problem and the research and measurement programs addressing it by:

8. Reference 1, page F-2.

9. W. J. Hill, P. N. Sheldon, J. J. Tiede, "Analyzing Worldwide Total Ozone for Trends," paper accepted for publication in Geophys. Res. Lett.

1. Providing a direct test of ozone depletion estimates from model calculations.
2. Maximizing the value and usefulness of the extensive ozone measurements, past, present, and future.
3. Providing a global early warning system for abnormal ozone changes, i.e., man-made or natural trends that are different from past variations.
4. Allowing practical limits to be set on the scope of the biological and climatic effects research programs in accordance with the maximum ozone change allowed by the early warning system plus appropriate action to limit the change.

To date the principal method used to test the ozone depletion theory has been indirect: measure the concentrations of certain species the amounts of which are quite small and compare the measurements with theoretical predictions. This has required the development of new data, techniques, and instruments, and the chlorofluorocarbon manufacturers have since 1974 had this as an important part of their research program. Among a number of projects relating to the analysis of ClO and other chlorine-containing species, one under the direction of Dr. D. H. Stedman at the University of Michigan measured successfully both ClO and Cl by titration with NO and detection of Cl by resonance fluorescence.¹⁰ NASA provided support for Dr. J. G. Anderson to develop a flight instrument based on this technique and make measurements of ClO and Cl this year. The values from the several sets of measurements vary several-fold from each other, and all are higher than theory. It will therefore be important to follow up this work in accordance with the NAS recommendations for:

1. Measurement of a greater number of different species simultaneously to provide a more rigorous test of the theory and
2. Measurement of the most important species simultaneously by two different techniques to check on the accuracy of the analytical methods.

10. D. H. Stedman, Res./Dev. 27, 22-4, 26, January, 1976.

Work supported by the chlorofluorocarbon manufacturers to help accomplish these tasks includes laboratory investigations of ultraviolet spectra of ClO by Dr. R. W. Nicholls of York University¹¹ and Dr. D. D. Davis of Georgia Tech and of the laser magnetic resonance technique for detection of ClO by Dr. C. J. Howard at the NOAA Laboratory at Boulder. Support is being provided for stratospheric measurements of ClO by Dr. D. G. Murcray using a balloon-borne ultraviolet spectrophotometer and to Dr. P. M. Solomon of SUNY Stony Brook for ground-based microwave measurements. In the latter program the 3 mm maser detector that will be employed will be the most sensitive receiver in the world for this type of observation.

To help resolve the serious questions that have been raised by the finding of larger than predicted amounts of chlorine species in the stratosphere, the chlorofluorocarbon manufacturers are supporting a project under Dr. A. E. J. Eggleton at AERE Harwell to develop methods for determining total chlorine and total fluorine. They are to be employed during balloon flights to compare their sum with the total amounts of these elements found with the sum of individual Cl- and F-containing molecules measured by other investigators.

3. "Develop an accurate materials balance--Inasmuch as a materials balance can detect, in principle, otherwise unidentified sinks for the CFMs, a detailed analysis should be made of how best to obtain a materials balance accurate enough to be meaningful. The study should define a minimum set of observations (including its time span) sufficient to establish a true global burden, and develop standards of sensitivity, calibration and accuracy necessary for the results to be useful. If such a study supports its feasibility, there should then be a coordinated observational program and data analysis for both F-11 and F-12."¹²

In spite of assertions to the contrary from elsewhere in the scientific community, the possibility of an unidentified sink

11. M. Mandelman and R. W. Nicholls, "The Absorption Cross-Sections and f-Values for the $v''=0$ Progression of Bands and Associated Continuum for the $\text{ClO}(A^2\Pi_1 + X^2\Pi_1)$ System," accepted for publication in J. Quant. Spectrosc. Radiat. Transfer.

12. Reference 1, pages F-2,3.

for the chlorofluorocarbons indicated above by the NAS Panel has been underscored by the NAS Committee:

"Although the materials balance can be interpreted as consistent with little or no active removal of the CFMs, the uncertainty limits range from zero inactive removal to a rate sixfold faster than that of the stratospheric photolysis."¹³

Modeling work sponsored by the chlorofluorocarbon manufacturers led to the realization that chlorofluorocarbon measurements to date did not rule out the possibility of sinks for these molecules.¹⁴ The most recent work indicates the most probable tropospheric lifetime for FC-11 is between 15-20 years.¹⁵ The ozone depletion calculations in the NAS reports are based on an infinite tropospheric lifetime and would be reduced as much as five-fold by this shorter tropospheric lifetime.

In its recommendation above the NAS Panel calls for development of both strategy and techniques to determine if the materials balance approach can be successful. The chlorofluorocarbon manufacturers have encouraged one of their modeling contractors to examine measurement problems and strategy. Several reports were prepared including one that was presented to the NASA Stratospheric Research Advisory Committee in December 1975 and has appeared subsequently as a publication.¹⁶

Three contractors in the industry program are now studying problems of calibration and accuracy of chlorofluorocarbon

13. Reference 2, page 5-10.

14. N. D. Sze and M. F. Wu, "Measurements of Fluorocarbons 11 and 12 and Model Validation: An Assessment," 27 pp., 15 December 1975.

15. J. P. Jesson, P. Meakin, and L. G. Glasgow, "The Fluorocarbon-Ozone Theory II: Tropospheric Lifetimes: An Experimental Estimate of the Tropospheric Lifetime of CCl_3F ," accepted for publication in Atmos. Environ.

16. R. G. Prinn, F. N. Alyea, and D. M. Cunnold, Bull. Am. Meteorol. Soc. 57, 686-94, June, 1976.

measurements, Drs. J. E. Lovelock, R. A. Rasmussen, and D. H. Stedman. Two are investigating new methods of absolute calibration, and one will conduct an interlaboratory comparison program.

4. "Reduce errors in rate constants--Considerable improvement in the rate constants for the $\text{Cl} + \text{CH}_4$ and $\text{HO} + \text{HO}_2$ reactions should be sought and may be expected within the next year or two. Further work on several of the other reactions in Table 8.3 would serve to reduce their lesser contributions to the uncertainties. A strong effort should be made to reduce the uncertainty factor of the $\text{HO} + \text{HO}_2$ reaction to 1.5 and those of the other reactions to 1.3, or less, including the effects of temperature dependences."

The uncertainty in the rate of the reaction $\text{HO} + \text{HO}_2$, of all the reactions in the model, accounts for the largest uncertainty in calculated ozone depletion. Because this is a reaction between two radicals, it is especially difficult to measure. The chlorofluorocarbon manufacturers are supporting two investigators to determine the rate of this and other reactions of the HO_2 radical. Drs. B. A. Thrush of the University of Cambridge and C. J. Howard of the NOAA-Boulder laboratory. Dr. J. W. Birks has measured the rate of the reaction $\text{ClO} + \text{NO}_2$, which is listed as another of the important reactions in the NAS Panel report.

5. "Detailed evaluation of identified sinks--As a complement to the materials balance approach, inactive removal processes of the CFMs, once identified, should be characterized in quantitative terms. In particular, oceanic removal of F-11 should be investigated more thoroughly by additional measurements of its concentration gradient in surface waters, by observations of its transport across the air-water interface, by redetermination of its solubility in sea water, and by a search for mechanisms that might contribute to its degradation in the surface waters. The same types of studies should be made for F-12. Also, efforts should be made to place narrower limits on the removal of F-11 and F-12 by photolysis and ion-molecule reactions in the troposphere."

A number of potential sinks for the chlorofluorocarbons have been investigated in the industry sponsored program including photochemical changes by Dr. C. Sandorfy of the University

17. Reference 1, page F-3.

18. Reference 1, page F-3.

of Montreal;¹⁹ reactions with plants and soil and with OH and the C(¹D) atom²⁰ by Dr. J. N. Pitts of the University of California at Riverside; reactions involving charged species by Dr. M. J. Campbell of Washington State University,²¹ Dr. V. A. Mohnen of SUNY Albany, and Dr. R. G. Hirsch of Du Pont;²² and trapping in Antarctic ice by Dr. R. A. Rasmussen. These projects have not identified with certainty a tropospheric sink for chlorofluorocarbons; studies by Dr. J. E. Lovelock are continuing.

C. "Improve other aspects of atmospheric chemistry--Photochemical processes and concentrations of natural species (NO_x and HO_x) contribute appreciable uncertainties to the prediction of ozone reduction by the CFMs. These aspects should be studied further, to place better limits upon their importance and to seek the most productive ways of reducing their contributions."²³

OH is among the HO_x species contributing to the uncertainties in the prediction of ozone depletion by the chlorofluorocarbons. The industry is supporting development of an instrument by Dr. D. D. Davis for measuring stratospheric OH concentrations by laser-induced fluorescence. A miniturized version of a unit flown successfully earlier on an airplane is being prepared for balloon flights, which will allow observations over extended periods of time, and of diurnal changes.

In accordance with the NAS Panel recommendation that important species be measured simultaneously by two different methods, Dr. D. G. Murcray is also being supported to measure the concentration of stratospheric OH by solar ultraviolet absorption.

19. C. Sandorfy, Atmos. Environ. 10, 343-51, May, 1976.

20. R. Atkinson, G. M. Brewer, J. N. Pitts, Jr., and H. L. Sandoval, "Tropospheric and Stratospheric Sinks for Halocarbons," accepted for publication in J. Geophys. Res.

21. M. J. Campbell, Geophys. Res. Lett. 3, 661-4, November, 1976.

22. R. G. Hirsch, Atmos. Environ. 10, 703-5, September, 1976.

23. Reference 1, pages F-3,4.

SUMMARY

Research Program on

EFFECT OF FLUOROCARBONS ON THE ATMOSPHERE

Sponsored by: The Fluorocarbon Industry

Prepared by: B. Peter Block
Frank A. Bower
Hillel Magid
J. Richard Soulen
Richard B. Ward

Distributed by: Manufacturing Chemists Association
1825 Connecticut Avenue, N. W.
Washington, D. C. 20009

(Originally Issued: September 26, 1975)

Revision No. 5: December 31, 1976

TABLE OF CONTENTS

	<u>Page</u>
The Industry-Sponsored Program	1
Gaps in Scientific Knowledge	5
Summary and Recommendations	7
 Tables:	
1 Fluorocarbon Manufacturers Represented on the MCA Technical Panel on Fluorocarbon Research	8
2 Fluorocarbon Research Program - Financial Summary	9
3 Fluorocarbon-Ozone Problem - Recommended Studies and MCA Program	10
4 Fluorocarbon Research Program - Types of Research Activities, Summaries	14
A. Reaction Rate Constant Measurements	14
B. Source and Sink Studies	15
C. Laboratory Studies Related to Potential Atmospheric Measurements	18
D. Tropospheric and Stratospheric Measurements	21
E. Modeling	24
F. Other	25
G. Consultants	26
5A MCA Projects - Work Completed	27
5B MCA Projects - Work in Progress	28
6 Publications from Work Supported by Fluorocarbon Manufacturers	30

SUMMARY

Research Program on EFFECT OF FLUOROCARBONS ON THE ATMOSPHERE

Sponsored by the Fluorocarbon Industry
Administered by the Manufacturing Chemists Association

(Originally Issued: September 26, 1975)
Revision No. 5: December 31, 1976

This summary describes work supported by the manufacturers of chlorofluorocarbons (commonly called fluorocarbons) in an attempt to assess the possible impact of these chemicals on the environment and, in particular, on the stratospheric ozone layer.

The Industry-Sponsored Program

In July of 1972, E. I. du Pont de Nemours & Company issued to fluorocarbon manufacturers worldwide an invitation to a "Seminar on the Ecology of Fluorocarbons." Its purpose was to establish a technical program because, as stated in the invitation, "Fluorocarbons are intentionally or accidentally vented to the atmosphere worldwide at a rate approaching one billion pounds per year. These compounds may be either accumulating in the

atmosphere or returning to the surface, land or sea, in the pure form or as decomposition products. Under any of these alternatives, it is prudent that we investigate any effects which the compounds may produce on plants or animals now or in the future."

Representatives of 15 companies attended the meeting, agreed that such a program was important, and established and funded a fluorocarbon research program under the administration of the Manufacturing Chemists Association (MCA). Thus, in 1972, with no evidence that fluorocarbons could harm the environment, the producers of these chemicals agreed unanimously that there was a need for more information and proceeded to act.

The 20 fluorocarbon producers supporting this program (see Table 1, p. 8) represent almost the total free world production of fluorocarbons. The research is directed by an MCA Technical Panel on Fluorocarbon Research with one member from each supporting company. This Technical Panel meets for two full days each month to review progress on current research, evaluate new proposals, and exchange data with contractors and with government agencies and researchers. Three projects were initiated in 1973:

1. Dr. James Lovelock, U. of Reading, "Fluorocarbons in the Environment."
2. Dr. Camile Sandorfy, U. of Montreal, "Spectroscopy and Photochemical Changes in Fluorocarbons."

3. Dr. O. C. Taylor, U. of California-Riverside,
"Monitoring and Atmospheric Reactions of Fluoro-
carbons."

Publication of the Rowland-Molina hypothesis in 1974 identified a potentially serious problem, so the MCA research program was expanded considerably. The fluorocarbon-ozone relationship attracted the attention of many scientists in academic and government laboratories, legislative and regulatory bodies, and the press. MCA's program is concentrating on research most likely to answer the critical question: to what extent will fluorocarbons affect the stratospheric ozone layer?

To strengthen the overall effort to find the answer, MCA has attempted to coordinate its efforts with others working on the same or related problems such as the SST and the space shuttle. All of these concern the federal government, and interactions with a number of agencies have been especially helpful in:

1. Taking advantage of the knowledge and experience gained in the Climatic Impact Assessment Program;
2. Coordinating funding of programs addressing the halogen-ozone problem;
3. Planning joint experiments with government research groups; and
4. Helping to set priorities for industry-sponsored research.

About 165 research proposals have been reviewed to date, and projects totaling over \$3,200,000 have been funded (see Table 2, p. 10). Calendar 1977 commitments are expected to exceed \$2,000,000, with total expenditures through 1977 expected to exceed \$5,000,000. Approximately \$650,000 has been committed to studies concerning the measurement of reactive chlorine species in the stratosphere, a difficult and important area. A promising method was developed at the University of Michigan with MCA support, and funding for the follow-on program for development of equipment and stratospheric measurements has been provided by the National Aeronautics and Space Administration (NASA). This illustrates an important contribution by industry: prompt funding of promising work that, if successful, can be taken over by a government agency for incorporation into its more comprehensive program.

MCA Technical Panel members have visited knowledgeable scientists throughout North America and Europe, including many at government laboratories. The panel held its September 1975 meeting in Montreal where, in addition to hearing presentations by Canadian contractors and potential contractors, it was briefed on the stratospheric measurement program of the Canadian government. The February 1976 panel meeting was held near Milan, Italy, and the October 1976 meeting in Erbach, Germany. The panel was briefed on pertinent European activities and heard presentations by European atmospheric scientists at these meetings.

Discussions in early 1975 with the Energy Research and Development Administration's (ERDA) Sandia Laboratory at Albuquerque and the Army's Atmospheric Sciences Laboratory at White

Sands Missile Range, New Mexico, indicated that industry could contribute significantly to the stratospheric composition (STRATCOM) program. Cooperation began immediately with the panel providing funding for two additional STRATCOM experiments and an additional balloon for stratospheric measurements in the fall of 1975. Similarly, the panel provided funding for several experiments that were flown on balloons included in the May 1976 Army COSMEP flights in Alaska and the STRATCOM flights in September 1976. In the 1976 flights there was also cooperation between MCA contractors and the Atmospheric Environment Service of Canada on Fourier transform spectrometer measurements. Through these types of collaboration it is hoped to reduce substantially the time required to obtain critical research results.

Gaps in Scientific Knowledge

Two groups were charged during 1975 with looking exclusively at the scientific aspects of the halocarbon-ozone problem and making recommendations for further work. In May 1975 the government's Interdepartmental Committee for Atmospheric Sciences (ICAS) made 12 recommendations for research and monitoring programs to assess the possible impact of fluorocarbons and halogens on ozone. Six of these were singled out as of major importance. In July 1975 the Panel on Atmospheric Chemistry of the National Academy of Sciences (NAS) identified a number of areas in which relevant data are nonexistent, fragmentary, or insufficient and listed 19 high priority areas requiring study. The final report of the NAS Panel, which was issued in September 1976, contains six recommendations for pertinent studies that are expected

to be completed in the period of up to two years that its parent committee, the NAS Committee on Impacts of Stratospheric Change, recommends be allowed before a decision is made on the necessity for restrictive action. The Panel also made fifteen recommendations for longer range studies, many of which are relevant to the possible impact of halogens on ozone.

Table 3 shows the relationship between the latest NAS recommendations and the research program sponsored by the fluorocarbon manufacturers (pp. 10 and 12) and with the preliminary recommendations of NAS and those of ICAS (pp. 11 and 13). There is generally good agreement between the ICAS and NAS recommendations. In addition to other small differences, the former include some aspects of monitoring not in the latter, and the latter include photolysis studies and bromine chemistry not in the former. The second column of Table 3 lists the MCA-administered industry program. Its emphasis is overwhelmingly in the major areas recommended by the NAS Panel and ICAS. The industry-sponsored program, therefore, aims to fill in the most important gaps in existing scientific knowledge. The only major area not in the MCA program is acceleration of ozone monitoring, which can be accomplished more appropriately by governmental and international agencies. In addition, the MCA program includes work to correlate UV radiation reaching the ground with ozone measurements and to improve the statistical treatment of ozone data so that very small abnormal changes can be detected in a much shorter period of time than was previously believed possible.

More detail on the MCA program is given in Tables 4, 5A, and 5B. Table 4 (p. 14) lists summaries of the projects by type of research activity. Table 5A (p. 27) lists completed projects, and Table 5B (p. 28) lists active projects in chronological order of funding. Table 6 (p. 30) lists publications resulting from industry sponsored work. These important contributions have not always been recognized as industry-funded activity.

In addition to the work supported by the industry as a whole there are also studies underway in the laboratories of individual member companies who have staff scientists able to make significant contributions to the resolution of the problem. Examples are the modeling, reaction, and spectroscopic studies at Du Pont and the statistical analysis of ozone trends at Allied.

Summary and Recommendations

In 1972 the fluorocarbon manufacturers began supporting a program to investigate the effects of fluorocarbons on the environment. This program has been expanded greatly to help determine the extent, if any, to which these compounds may affect the stratospheric ozone layer. Industry and government-sponsored scientists working on the halogen-ozone problem have cooperated effectively. Continuation of this cooperation is essential, with special attention to providing periodically updated summaries of research priorities, programs, and results, together with critical analyses of the reliability and significance of the data. Such a program will produce most rapidly the information required for rational decision making.

Table 1

FLUOROCARBON MANUFACTURERS

represented on the

MCA TECHNICAL PANEL ON FLUOROCARBON RESEARCH

Akzo Chemie nv (Holland)
Allied Chemical Corporation (U.S.)
Asahi Glass Co., Ltd. (Japan)
Australian Fluorine Chemical Pty. Ltd. (Australia)
Daikin Kogyo Co., Ltd. (Japan)
E. I. du Pont de Nemours & Company, Inc. (U.S.)
Du Pont of Canada Limited (Canada)
Hoechst AG (West Germany)
Imperial Chemical Industries Limited (England)
I.S.C. Chemicals Ltd. (England)
Kaiser Aluminum & Chemical Corporation (U.S.)
Kali-Chemie Aktiengesellschaft (West Germany)
Mitsui Fluorochemicals Co. Ltd. (Japan)
Montedison S.p.A. (Italy)
Pennwalt Corporation (U.S.)
Racon Incorporated (U.S.)
Rhône-Poulenc Industrie (France)
Showa Denko K. K. (Japan)
Ugine Kuhlmann, Produits Chimiques (France)
Union Carbide Corporation (U.S.)

December 31, 1976

Table 2

FLUOROCARBON RESEARCH PROGRAM

Administered by
Manufacturing Chemists Association

FINANCIAL SUMMARY

	Completed Projects	Active Projects	Total
Chemical Reactions and Kinetics	\$ 242,050	\$ 351,600	\$ 593,650
Analytical Methods	259,312	105,418	364,730
Tropospheric Measurement	169,107	339,846	508,953
Stratospheric Measurement	211,187	1,036,530	1,247,717
Modeling	147,652	285,005	432,657
Consulting	<u>31,475</u>	<u>22,646</u>	<u>54,121</u>
TOTAL	\$1,060,783	\$2,141,045	\$3,201,828

December 31, 1976

Table 3

FLUOROCARBON-OZONE PROBLEM - RECOMMENDED STUDIES AND MCA PROGRAM

NAS Recommendations, September 1976 ^{a,b}	MCA Supported Work	
	Completed	In Progress
I.1. Establish role of ClONO ₂ in the stratosphere.		Birks - U. of Illinois Murray - U. of Denver Timmons - Catholic Wiesenfeld - Cornell
I.2. Verify the reliability of predicted ozone reductions. a. via ClO and Cl profiles simultaneously with HCl, O ₃ , etc. b. via decrease in total ozone column c. via changes in ozone profile	Davis - U. of Maryland ^f Ekstrom - Battelle Northwest Stedman - U. of Michigan ^f Young - Xonics, Inc. ^f	Davis - U. of Maryland Institute of Techn Howard - NOAA-Boulder Murray - U. of Denver Nicholls - York U. ^f Parzen and Pagano - Science and Techn Foundation, Inc. Solomon - State U. of Stony Brook
I.3. Develop an accurate materials balance for F-11 and F-12. a. Establish requirements for sensitivity, calibration, and accuracy. b. Define minimum observational program.	Lovelock - U. of Reading Manufacturers via MCA Rasmussen - Washington State U. Taylor - U. of California-Riverside	FARLTON - AERE Bar Lovelock - Private Murray - U. of Denver Manufacturers via MCA Rasmussen - Washington Stedman - U. of Mich
I.4. Reduce errors in rate constants with emphasis on Cl + CH ₄ , OH + HO ₂ , ClO + O, ClO + NO, ClO + NO ₂ , OH + HCl, Cl + O ₃		Birks - U. of Illinois Howard - NOAA-Boulder Thrush - U. of Cambr Timmons - Catholic Wiesenfeld - Cornell
I.5. Make a detailed evaluation of sinks once they are identified. a. Oceanic removal of F-11 and F-12. b. Photolysis of F-11 and F-12. c. Ion-molecule reactions of F-11 and F-12 in the troposphere.	Campbell - Washington State U. Mohnen - State U. of N.Y.-Albany Pitts - U. of California-Riverside	Birks - U. of Illinois Kaufman - Emory U. Lovelock - U. of Reading Martin - Aerospace Rasmussen - Rasmussen
I.6. Improve other aspects of atmospheric chemistry. a. Photochemical processes of natural species (NO _x , HO _x). b. Concentrations of natural species.		Bulls - Bomen, Inc. Davis - U. of Maryland Institute of Techn Girard - ONERA (France) Murray - U. of Denver
II.1. Monitor all major releases of volatile halocarbons and other potential stratospheric pollutants on a global scale.	Lovelock - U. of Reading Rasmussen - Washington State U. Taylor - U. of California-Riverside	Lovelock - Private Murray - U. of Denver Rasmussen - Washington Kidley - York U.
II.2. Determine sinks for every substance identified as a potential or actual hazard to the stratosphere.		cf. I.5
II.3. Determine for each pollutant the complete sequence of events from injection to removal of the final products from the atmosphere. a. Fate of halocarbon radicals remaining after photodissociation. b. Effect of HF and HCl on rain and snow.		
II.4. Improve reliability of 1-D calculations. a. More extensive and more accurate data for CH ₄ , N ₂ O, HF, and O ₃ for "fine-tuning" models. b. Treatment of diurnal variations.		Cunnold, Alveda, Princeton ates Sze - ERT, Inc.
II.5. Expedite development of 2- and 3-D models.		Cunnold, Alveda, Princeton ates
II.6. Determine accurate rate constants for a large number of chemical reactions involving ClX and atmospheric species, HO _x chemistry, and O(1D) quenching and reaction rate constants.		cf. I.4
II.7. Determine reaction rate parameters over the tropospheric and stratospheric temperature range for the reaction of OH with all important chloro- and fluorocarbons.		

1. Measure Cl and ClO concentrations in the stratosphere.

1. Measure Cl and ClO upper atmosphere concentration profiles.

5. Accelerate program for monitoring of ozone on a global basis. Establish trends in atmospheric ozone.

3. Make systematic search for other chlorine-bearing gases in the stratosphere.

14. Determine total chlorine in the atmosphere and compare with sum of all identified compounds.

A2. Expand monitoring of F-11, F-12, and CCl₄ in the troposphere and develop standards for their measurement.

6. Confirm and improve certain reaction rate measurements crucial to ozone destruction.

6. Measure rate constants for ClO + SO, ClO + O, Cl + HO₂, Cl + CH₄.

7. Measure rate constants for OH + H₂O; H, O, NO, C₃ + HO₂; H₂O, CH₄, N₂, O₂ + O(1D).

4. Conduct promising research to determine sinks for chlorine in the stratosphere.

16. Seek out and characterize other possible sinks for removal of chlorofluorocarbons.

17. Determine source and further reactions of chlorine in stratospheric particulates.

2. Measure [F-11, F-12, CCl₄, OH, H₂O, and other gases at 25-40 km.

2. Measure OH, HCl, HF, NO, etc., upper atmosphere concentration profiles.

12. Sample at low altitude to find additional possible natural and anthropogenic sources of halogen-containing compounds. Establish concentrations.

2. Measure F-11, F-12, CCl₄, [OH, H₂O,] and other gases at 25-40 km.

4. Measure F-11, F-12, CCl₄, CHCl₃, and N₂O concentration profiles 0-40 km. Establish standards.

A1. Obtain a full record of past and present halocarbon production and release.

13. Obtain accurate data on production and uses of each halocarbon manufactured worldwide.

Cf. I.5

Cf. I.5

A3. Study the lifetimes of halocarbons, HCl, and phosgene in the troposphere.

15. Measure atmospheric abundance of decomposition products such as COF₂, COCl₂, COFCl, and determine whether COFCl releases additional Cl.

A4. Expand the stratospheric modeling studies of ozone formation and destruction. Improve chemistry and transport.

3. Measure global distribution of certain trace gases to improve models.

5. Develop better atmospheric models.

Cf. II.4

Cf. I.4

Cf. I.4

Cf. I.4

8. Measure rate constants for OH + CH₂FCl, CHF₂Cl, CH₃Cl, CH₃CCl₃, CH₂Cl₂, and all other potentially usable halocarbons with C-H bonds.

Table 1 (continued)

MCA Supported Work

NAS Recommendations, September 1976^{a,b}

	Completed	In progress
II.8. Determine kinetic data for all important reactions of BrX and photolytic properties of the various bromine-containing species.		
II.9. Measure solar flux and its variability, especially at wavelengths greater than 175 nm, and stratospheric scattering parameters.		
II.10. Study photochemistry of halocarbons and CIX further. a. Photolysis parameters of F-11, F-12, CClO, CClO ₂ , and ClONO ₂ . b. Photolysis parameters of other halocarbons such as CHFCI ₂ , CCl ₄ , CH ₃ CCl ₃ , CH ₂ Cl ₂ , and CH ₃ Cl.	Sandorfy - U. of Montreal	
II.11. Define and establish a global network for atmospheric measurements.	Ekstrom - Battelle Northwest	Solomon - State U. of N. Y. - State Brook
II.12. Measure concentration profiles in upper atmosphere for species related to the chemistry of chlorine, including OH, HCl, HF, and NO.		Butts - Boeing, Inc. Davis - U. of Maryland/Georgia Institute of Technology Girard - ONERA (France) ⁶ Murray - U. of Denver
II.13. Make global monitoring of ozone sufficiently sensitive, accurate, and complete to detect small changes due to man's activities.		Parzen and Pagano - Frontier Science and Technology Research Foundation, Inc.
II.14. Give further attention to the problem of water vapor feedback.		
II.15. Study the reduction in stratospheric ozone associated with the use of nitrogen fertilizers including the effects of the interaction between the NO _x and ClO _x cycles.		Connolly, Alcoa, Prinn - (AF Associates)
	Nicholls - York U.	Davis - U. of Maryland/Georgia Institute of Technology
a. "Halocarbons: Effects on Stratospheric Ozone," Panel on Atmospheric Chemistry, Committee on Impacts on Stratospheric Change, Assembly of Mathematical and Physical Sciences, National Research Council, NAS, September, 1976, Appendix F.		
b. Group I items are short-term (1-2 year), Group II longer range.		
c. "The Possible Impact of Fluorocarbons and Halocarbons on Ozone," ICAS 18a-FY 75, May, 1975, pp. 65-66.		
d. 1-6 indicate major recommendations, A1-A6 other.		
e. "Interim Report of the Panel on Atmospheric Chemistry," Climatic Impact Committee, Assembly of Mathematical and Physical Sciences, National Research Council, NAS, July, 1975, pp. 1-8.		
f. Laboratory study to develop data, analytical method, or instrumentation required for stratospheric measurements.		
g. Office National d'Etudes et de Recherches Aérospatiales.		

S Recommendations, May 1975^{c,d}

NAS Recommendations, July 1975^e

19. Obtain kinetic data for all important reactions of bromine, $\text{Br} + \text{O}_3$, HO_2 ; $\text{BrO} + \text{O}$, NO , O_3 ; $\text{CH}_3\text{Br} + \text{OH}$.

9. Measure solar flux and scattering parameters for the stratosphere.

10. Measure photolysis parameters for chlorohydrocarbons and chlorofluorocarbons.

Expand monitoring of F-11, F-12, and CCl_4 in the troposphere (and develop standards for their measurement).

18. Accurate baseline monitoring of ground level composition and absorption spectrum of "clean air" including measurements of O_3 , HCl , HF , N_2O , NO , NO_2 ; monitoring of column densities of HCl , HF , ClO , OH , NO_2 , O_3 from ground level and satellite stations.

I.6 and II.1

Cf. I.6, II.11

I.2

Cf. I.2

Expand the general circulation modeling of the effects of reduced stratospheric ozone in climate.

Continue to study the effects of releasing halocarbons on humans and the biosphere.

11. Make laboratory studies of the photolysis of ClO .

December 11, 1976

Table 4
Fluorocarbon Research Program
Types of Research Activities, Summaries

A. Reaction Rate Constant Measurements

Dr. J. W. BIRKS -- University of Illinois -- 75-1, 76-117A. Measurement of Reaction Rates Relevant to the Fluorocarbon-Ozone Problem.

Reaction rates are measured at various temperatures by a discharge flow technique with a quadrupole mass spectrometer for detection of products. Reactions studied include $\text{ClO} + \text{O}_3 + \text{M}$ (yields OClO , unimportant in daytime), $\text{ClO} + \text{NO}_2 + \text{M}$ (important in lower stratosphere, $k = 1.5 \times 10^{-13} \text{cm}^6 \text{ molecule}^{-2} \text{s}^{-1}$ at 297°K), $\text{ClO} + \text{SO}_2$ (no appreciable reaction observed), $\text{NO} + \text{O}_3$ (in agreement with previous studies), $\text{O} + \text{ClONO}_2$ (preliminary results in agreement with Davis), and $\text{BrO} + \text{NO}_2$ (in progress). The products of the four-center reactions $\text{Cl}_2\text{O} + \text{Cl}$, O , N , or ClO have been investigated by molecular beam mass spectrometry and indicate that O , N , and ClO all react with the O in ClOCl . Future work will include $\text{ClO} + \text{H}_2\text{O}_2$ and reactions of HO_2 .

Dr. C. J. HOWARD -- NOAA Boulder -- 76-100. Laser Magnetic Resonance Study of HO_2 Chemistry.

The object of this study, which is just getting under way, is to use the laser magnetic resonance technique to measure rate constants for stratospherically important HO_2 reactions.

Dr. J. N. PITTS -- University of California at Riverside -- 74-2. Atmospheric Reactions of Fluorocarbons.

Reaction rate constants have been measured for the reactions of $\text{O}(^1\text{D})$ with fluorocarbons 11, 12, 22, 113, and 114, and of OH with fluorocarbons 11, 12, and 22. The results indicate that in the stratosphere the reaction of $\text{O}(^1\text{D})$ atoms with fluorocarbons 11 and 12 is secondary to photolysis, whereas the reaction of OH with fluorocarbon 22 is much more important than photolysis. The photooxidation products of 12, 13, and 22 at 184.9 nm, i.e., COCl_2 , COFCl , and COF_2 as appropriate, are also observed to be the products for reaction with $\text{O}(^1\text{D})$.

Table 4 (continued)

Dr. B. A. THRUSH -- University of Cambridge -- 75-58, 75-58II. Reactions of the HO₂ Radical Studied by Laser Magnetic Resonance.

The rate coefficient of the reaction O + HO₂ has been measured for the first time. The value found, $4.3 \times 10^{-11} \text{ cm}^3\text{s}^{-1}$, improves the fit of the calculated OH profile with Anderson's recent measurements. Work on the HO + HO₂ reaction is in progress.

Dr. R. B. TIMMONS -- Catholic University -- 76-129. Photochemical and Chemical Kinetics Measurements of Stratospheric Importance With Respect to the Fluorocarbon Issue.

It is planned to measure accurately the UV absorption cross-section of HOCl, study its photochemistry, and perform ClONO₂ photochemical studies to complement Wiesenfeld's proposal (q.v., below). HOCl is a possible stratospheric sink the magnitude of which would depend on the absorption cross-section. Earlier spectral measurements were inaccurate.

Dr. J. WIESENFELD -- Cornell -- 76-128. Photochemistry of Small Chlorinated Molecules.

The photochemistry of chlorine nitrate will be studied. While not yet thoroughly treated in the models, chlorine nitrate may be an important stratospheric sink. A flash photolysis-resonance absorption system is used for photolysis and analysis of products and incorporates either a conventional flash lamp with filters or a tunable laser. The wavelength and temperature dependencies of the quantum yield and product formation will be determined.

B. Source and Sink Studies

Dr. J. W. BIRKS -- University of Illinois -- 76-117B. Studies of Heterogeneous Reactions.

Potentially important heterogeneous reactions, which will be chosen with the advice of ERT, are to be studied, including the heterogeneous reaction of ClONO₂ and H₂O. Equipment is being assembled for measurements in early 1977.

Table 4 (continued)

Dr. M. J. CAMPBELL -- Washington State University -- 75-53. Chlorofluoromethane Destruction by Natural Ionization.

Laboratory measurements under conditions approximating the real atmosphere in all respects except ion age show large rate constants for removal of CCl_4 and fluorocarbon 11, apparently by atmospheric ions. The rate constant for fluorocarbon 12 is much smaller. If the same rate constants were to hold for older ions, this mechanism might constitute a sink that could decrease the stratospheric concentration of fluorocarbon 11 by a factor of two.

Dr. M. KAUFMAN -- Emory University -- 76-126. Studies of Compounds of Sulfur, Oxygen, and Chlorine.

Reactions between SO_x and Cl_x species of possible stratospheric importance are being investigated. They may yield compounds with long lifetimes or lead to incorporation of Cl into aerosols. Currently, the rate constant of the reaction of Cl with SO_2 is being measured.

Dr. J. E. LOVELOCK -- University of Reading -- 75-67. Unidentified Factors in the Fluorocarbon-Ozone Problem.

The absorption and reactivity characteristics of the atmospheric dust plume from the Sahara for atmospheric halocarbon vapors (fluorocarbon 11 and CCl_4) are being evaluated. Investigations are in progress on the relationship between photochemically produced atmospheric peroxy compounds (e.g., peroxyacetyl nitrate) and the incidence of skin carcinoma.

Dr. L. R. MARTIN -- Aerospace Corp. -- 75-81. Laboratory Investigation of the Heterogeneous Interaction of Cl and ClO With H_2SO_4 .

A flowing afterglow apparatus is being used to measure the rate of the heterogeneous reactions of Cl and ClO with sulfuric acid, simulating the stratospheric aerosol. Preliminary data suggest that the reaction rate of Cl is too slow for its reaction to constitute a significant sink, although rates were markedly increased by the presence of certain metal salts in the sulfuric acid. Measurements of the ClO rates, to which the ozone depletion cycle is more sensitive, are planned. The nature of the surface interaction is also to be characterized.

Table 4 (continued)

Dr. V. A. MOHNEN -- SUNY Albany -- 75-64. Ion Molecule Reactions Involving Fluorocarbons.

A drift tube in which ions are formed and their mobility determined and an electric quadrupole mass spectrometer were used to study the role of ion-molecules in the reactions between fluorocarbon 12 and positive and negative ion clusters that form in air-like gas mixtures. There was no clustering of fluorocarbon 12 and no evidence for charge transfer or cluster ion reactions with positive ions of the form $H^+ \cdot (H_2O)_n (NH_3)_m$. Reaction with negative ions of the form $O^- \cdot X$ and $CO_4^- \cdot X$ results in the formation of the ion-cluster series $Cl^- \cdot (H_2O)_x$.

Dr. J. N. PITTS -- University of California at Riverside -- 75-12. Monitoring and Atmospheric Reactions of Fluorocarbons.

Long-term photolysis studies in simulated sunlight revealed that fluorocarbons 11 and 12 are photochemically stable, even when irradiated for several weeks. An attempt to study fluorocarbon- NO_x -air mixtures under simulated stratospheric conditions could not be completed because the background level of hydrocarbons in the environmental chamber could not be made sufficiently small.

Plant tissues did not absorb measurable quantities of fluorocarbons 11, 12, or 22, and no adverse effects could be measured. Fluorocarbons penetrate into the soil atmosphere, and concentrations change in direct relationship with changes in concentration in the atmosphere above ground.

Dr. R. A. RASMUSSEN -- Washington State University -- 75-71. Measurement of Fluorocarbon Content of "Antique" Air Samples.

A sensitive method for the determination of low parts per trillion analysis of fluorocarbons 11 and 12 in small-volume air samples in containers was developed and applied to a wide variety of vessels believed to contain antique air. Both of these fluorocarbons were found in all samples analyzed, suggesting either the presence of natural fluorocarbons before 1930 or contamination during storage or handling.

Table 4 (continued)

Dr. R. A. RASMUSSEN -- Washington State University --
75-84. Collection and Analysis of Antarctic Ice Cores.

The concentration of halocarbons in air obtained from Antarctic snow shows no enrichment in samples obtained from the Ross ice shelf (mainly -30°F), whereas there is enrichment in samples obtained from the South Pole (-50 to -60°F).

Dr. R. A. RASMUSSEN -- Private -- 76-140. Lower Stratospheric Measurement of Non-methane Hydrocarbons.

This study is now beginning. It is important because non-methane hydrocarbons, if present in the stratosphere, will act as a sink for Cl in the same way that methane acts and thus reduce the ozone depletion attributable to chlorine-containing species.

Dr. C. SANDORFY -- University of Montreal -- 73-2. Spectroscopy and Photochemical Changes of Fluorocarbons.

The vacuum ultraviolet and photoelectron spectra of all industrially important chlorofluorocarbons were measured and published. The photochemical vulnerability of these molecules was predicted from their spectra.

C. Laboratory Studies Related to Potential Atmospheric Measurements

Dr. H. J. BUIJS -- Bomem, Inc. -- 75-90. Construction of a Fourier Transform Spectrometer.

A spectrometer with a projected resolution of 0.01 cm^{-1} is being completed for use in the simultaneous determination of ClONO_2 and either HCl or HF, or of HCl and HF.

Dr. D. D. DAVIS -- University of Maryland -- 74-10. Laboratory Determination of the Sensitivity of Laser-Induced Fluorescence for the Detection of ClO Under Atmospheric Conditions.

Ground-state stationary ClO concentrations of about 10^{12} cm^{-3} were scanned at several electronic absorption wavelength regions with a tunable UV laser. Laser-induced fluorescence proved to be unusable for measuring ClO because (1) at 300°K the highest populated K level, 12, contains only 5% of the total ClO and (2) the majority of the ClO undergoes predissociation to Cl and O.

Table 4 (continued)

Dr. D. D. DAVIS -- University of Maryland/Georgia Institute of Technology -- 75-73. Laboratory Measurement of Spectroscopic Absorption Cross-Sections of ClO.

Work is being completed on the measurement of the absorption cross-sections for ClO, data required for determining concentrations from UV absorption measurements of this molecule.

Dr. D. D. DAVIS -- University of Maryland/Georgia Institute of Technology -- 75-87. Development of Instrument for Stratospheric OH Measurement by Laser-Induced Fluorescence.

This program is to measure stratospheric OH concentrations over extended periods of time, including some covering diurnal changes. A miniaturized dye laser module for balloon flights has been built and tested. An engineering flight is planned after a high-altitude, pressurized chamber for housing the dye laser and Nd-Yag driver has been built and required data storage and control facilities are ready.

Dr. A. E. J. EGGLETON -- AERE Harwell -- 76-116. Total Chlorine Measurements in the Troposphere and Stratosphere.

The feasibility of measuring total chlorine and fluorine in the atmosphere by neutron activation and γ photon activation, respectively, after collection of reactive species and particulate material on a filter and collection of gaseous compounds on activated charcoal is being studied.

Dr. C. J. HOWARD -- NOAA, Boulder -- 75-47. Laboratory Determination of the Feasibility of Laser Magnetic Resonance for ClO Detection and Reaction Studies.

This study is to detect ClO using laser magnetic resonance and to utilize this method for measuring rate constants for stratospherically significant ClO reactions. It has been demonstrated that ClO can be detected by this method, and the sensitivity is under evaluation.

Dr. J. E. LOVELOCK -- Private -- 76-120. The Electron Capture Detector as a Reference Standard for Analysis of Atmospheric Halocarbons.

The electron capture gas chromatograph is being investigated as a reference standard for analysis of atmospheric halocarbons.

Table 4 (continued)

Dr. D. G. MURCRAY -- University of Denver -- 75-92.
Laboratory Measurement of High Resolution Infrared Spectra
of Chlorine-Containing Molecules of Stratospheric Interest.

Laboratory measurements of the high resolution infrared spectra of chlorine-containing molecules are to be made. Statistical-band-model analyses and integrated intensity measurements for the 10.8 μm band of fluorocarbon 12 and 11.8 μm band of fluorocarbon 11 have been published. Additional spectra for methane, methyl chloride, and carbonyl fluoride have been run, and work is in progress on chlorine nitrate.

Dr. R. W. NICHOLLS -- York University -- 75-11 and 75-11-II. Experimental and Theoretical Studies on the UV Spectrum of ClO With Stratospheric Applications.

Absolute absorption coefficients and cross-sections have been measured for all bands and the photodissociation continuum of the $V''=0$ progression for ClO. The very complicated emission spectrum that has been excited over the wavelength range 2500-4500 Å in discharges through ClO₂ and Cl₂O is currently undergoing measurement, identification, and analysis. Computer-based synthetic spectra of various ClO bands have been calculated. Emphasis is now being placed on the high resolution spectrum of ClO in the spectral region above 3000 Å, which should lay the groundwork for the detection and measurement of stratospheric ClO from the ground and from balloon-borne equipment. Possible interferences by O₃ are being evaluated.

Dr. R. W. NICHOLLS -- York University -- 75-30b.
Laboratory Studies of the Infrared Vibration-Rotation Spectrum of ClO.

Work in this area was suspended to allow greater effort in the UV measurements (75-11).

Dr. R. A. RASMUSSEN -- Private -- 76-142. Interlaboratory Comparison of Fluorocarbon Measurements.

A second round of identical samples of rural air will be circulated blind to participating laboratories for analysis for fluorocarbons 11 and 12, chloroform, methyl chloroform, carbon tetrachloride, and N₂O. The results will be analyzed statistically.

Table 4 (continued)

Dr. D. H. STEDMAN -- University of Michigan -- 74-7.
Atmospheric Determination of ClO Concentration: A Feasibility Study.

Laboratory studies have demonstrated the feasibility for detecting stratospheric ClO by chemical conversion to Cl (by reaction with NO) accompanied by vacuum ultraviolet resonance fluorescence. In-flight use of this technique is being supported by NASA.

Dr. D. H. STEDMAN -- University of Michigan -- 76-132.
Absolute Calibration of Fluorocarbon Measurements.

The final design of a feed-back flow system for the calibration of fluorocarbon samples is being built.

D. Tropospheric and Stratospheric Measurements

Dr. H. L. BUIJS -- Bomem, Inc. -- 75-98. Measurement of HCl and HF in the Stratosphere by Fourier Transform Spectroscopy.

Data have been successfully collected from balloon flights in Alaska (May 1976) and New Mexico (September 1976) and are being analyzed for HCl and HF concentrations.

Dr. P. A. EKSTROM -- Battelle Memorial Institute, Pacific Northwest Laboratories -- 75-27. Ground-Based Millimeter-Wavelength Observations of Stratospheric ClO.

About 500,000 data points were obtained in the microwave spectra near 93 GHz with the Kitt Peak radiotelescope. Excessive noise made interpretation difficult, but base-line corrected spectra suggested an upper limit on ClO of one hundred times model predictions.

Dr. A. GIRARD -- ONERA - France -- 75-88. Measurement of HCl, HF, ClO, etc., in the Stratosphere by High Resolution Infrared Spectroscopy.

Balloon-borne flights and supporting laboratory work are planned to measure fluorocarbons, HCl, HF, ClO, ClNO₃, and other compounds by high resolution infrared spectroscopy. This program is part of a cooperative program including aircraft flights with the Centre National d'Etudes Spatiales. A flight in 1976 provided an HCl concentration profile between 25 and 30 km. Equipment failure occurred in a September 1976 flight. A flight from Brazil in December 1976 and additional flights in the spring of 1977 are planned.

Table 4 (continued)

Dr. J. E. LOVELOCK -- University of Reading -- 73-1, 74-3, 75-67. Fluorocarbons in the Environment.

The electron-capture gas chromatograph (ECGC) has been developed and applied to the measurement of chlorocarbons and chlorofluorocarbons in the troposphere and lower stratosphere, particularly over Europe and the Atlantic Ocean. Current fluorocarbon 11 levels are about 130 ppt (U.K.) and 80 ppt (Southern Hemisphere). ECGC has been adapted to measure CH_3Cl at atmospheric concentrations of about 10^{-9} v/v, with sources for CH_3Cl identified as the ocean and smouldering vegetation.

Dr. D. G. MURCRAY -- University of Denver -- 75-13, 76-101, 76-135. The Measurement of the Stratospheric Distribution of Fluorocarbons and Other Constituents of Interest in the Possible Effect of Chlorine Pollutants on the Ozone Layer.

Measurement of stratospheric distribution by balloon-borne, high-resolution, infrared absorption measurement at large solar zenith angles has been achieved using a specially-constructed grating spectrometer. The distributions for fluorocarbon 11, fluorocarbon 12, and CCl_4 obtained show a concentration increase of about 2.5, with a rather wide range of uncertainty, from 1968 to 1975 for fluorocarbon 11 and fluorocarbon 12. Subsequent flights also yielded HCl and HNO_3 profiles. Profiles or upper limit measurements on several other species, including ClONO_2 , are potentially accessible.

Balloon-borne experiments will be continued to measure stratospheric ClO (300-500 nm) and OH (306.4 nm) by solar UV absorption.

Dr. R. A. RASMUSSEN -- Washington State University -- 75-2, 75-59. Fluorocarbon Research.

An attempt was made to obtain halocarbon concentration measurements as far into the stratosphere as could be reached by an available commercial aircraft. A small portable gas chromatograph was used for on-board measurements, and cannister samples were collected for subsequent detailed halocarbon analysis on the ground. One phase of the study consisted of samples collected over a wide area of the Pacific Northwest, a second of samples collected frequently to the maximum attainable altitude over Alaska. The halocarbon

Table 4 (continued)

concentrations are either constant or decrease very slowly with altitude in the troposphere, decrease rapidly in the tropopause from the tropospheric concentration to an average value identifiable with the stratosphere, and do not show a clear pattern of concentration gradients above the tropopause.

A trans-Pacific flight from 80°N to 60°S has been completed, and the air samples collected have been analyzed for halocarbons. Most of the samples were collected at 39,000 to 43,000 ft. Fluorocarbon 12 concentrations are about 10% higher in the north than in the south at ground level, and the difference is apparently greater for fluorocarbon 11.

Dr. B. A. RIDLEY -- York University -- 76-102A, 76-102B. Measurement of Fluorocarbons and Related Chlorocarbons in the Stratosphere by Collection and Analysis.

Balloon-borne grab sample devices have been developed and flown to obtain fluorocarbon 11, fluorocarbon 12, and N₂O concentrations at 10-40 km altitude. The data show a relatively consistent drop-off with height above the tropopause, whereas their consistency in drop-off is much worse as a function of height above the surface. The spread in data is larger than the accuracy of the measurements, which indicates variability in stratospheric concentrations due to horizontal motions.

Dr. P. M. SOLOMON -- SUNY Stony Brook -- 76-130. Millimeter Wave Observations of Chlorofluoromethane Byproducts in the Stratosphere.

This study, which is directed toward the development of a ground-based method for continuous determination of ClO, is just getting under way. One of the millimeter wave observing systems will be based on a 3 mm maser, a unique instrument that is the most sensitive detector in the world in the 83-94 GHz range.

Dr. O. C. TAYLOR -- University of California at Riverside -- 73-3, 74-2. Monitoring and Atmospheric Reactions of Fluorocarbons.

An electron capture gas chromatograph was used to measure the concentrations of fluorocarbons 11 and 12 in the troposphere over southern California and in the lower stratosphere over New Mexico and Colorado. The tropospheric concentrations were found to

Table 4 (continued)

vary from day to day as climatic conditions affected dispersion and dilution. Concentration decreased with increasing altitude in the lower stratosphere.

Dr. R. A. YOUNG -- Xonics, Inc. -- 75-50, 75-86.
Development of an Instrument to Measure O, ClO, O₃, and Total Cl in the Stratosphere.

The preliminary experiment to measure stratospheric ClO and total chlorine by resonance fluorescence flown on the September 1975 STRATCOM balloon was not completely successful; however, several key components appeared to have functioned satisfactorily. Follow-up work has been undertaken in a stratospheric simulator to provide design information for further development of stratospheric instrumentation.

E. Modeling

Drs. D. M. CUNNOLD, F. N. ALYEA, and R. G. PRINN -- CAP Associates -- 75-24, 76-122, 76-122S. Meteorological and Multi-Dimensional Modeling Considerations Relating to Atmospheric Effects of Halocarbons.

Studies to assess the accuracy and shortcomings of the 1-D model used to estimate ozone depletion indicated that a tropospheric lifetime for fluorocarbons 11 and 12 as short as 10 years was not inconsistent with atmospheric measurements. Thus, ozone depletion estimates might be considerably less than present estimates. The great variability of stratospheric measurements indicates that simultaneous measurements for many important stratospheric species are needed and that seasonal dependence of species must be considered.

The absence of circulation in the radiative models used to calculate warming (greenhouse effect) limits the value of calculated effects.

Because it has been shown that meteorological conditions can cause variations in both tropospheric and lower stratospheric fluorocarbon measurements in Alaska, meteorological analysis will be performed for selected fluorocarbon measurements, and temporal and latitudinal variability of fluorocarbon concentration will be determined using 2-D and 3-D models.

Table 4 (continued)

Dr. N. D. SZE -- Environmental Research and Technology, Inc. -- 75-32, 76-115. Model Analysis of the Fluorocarbon Problem.

A one-dimensional model has been used to evaluate the role of stratospheric water in the NO_x and ClX cycles, the relationship of eddy diffusion coefficient and fluorocarbon lifetime, the use of fluorocarbon measurements to calculate lifetime, and the effect of chlorine nitrate. The importance of OH concentration on calculated ozone depletion was shown, and key reactions were identified. Analyses showed 10-20 year tropospheric lifetimes were not inconsistent with measurements and helped to define quantitatively the uncertainties associated with ozone depletion calculations. Current work is to continue a sensitivity analysis of the 1-D model, develop a diurnal 1-D model, and include multiple scattering and heterogeneous (aerosol) reactions in the model.

F. Other

Dr. D. BERGER - Temple University -- 75-62. Ground-Level Monitoring of Ultraviolet Solar Radiation.

The monitoring of solar ultraviolet radiation, initiated by the Climatic Impact Assessment Program (CIAP), is being continued. The number of instruments and stations is being increased. Calibration of operating instruments is maintained, and comparisons between different types of monitoring equipment are being made.

Drs. E. PARZEN and M. PAGANO -- Frontier Science and Technology Research Foundation, Inc. -- 76-106. Total World Ozone Level: Statistical Analysis.

Ozone column measurements from at least 20 stations are being evaluated statistically to detect trends in recorded ozone concentrations and to establish the limits of detection for such trends. Time series analysis has been shown to be substantially more sensitive in detecting non-random ozone changes than the estimates of such sensitivity made by the Federal Task Force on Inadvertent Modification of the Stratosphere (IMOS) in 1975. The absence of detectable trends provides an upper limit for actual depletion and a test of model predictions.

Table 4 (continued)

If sufficient sensitivity is achieved, this technique will enable an effective early warning system for ozone depletion to be established.

G. Con ants

Dr. J. G. Anderson	University of Michigan	Stratospheric Measurements
Dr. A. W. Castleman, Jr.	University of Colorado	Heterogeneous Chemistry
Dr. F. C. Fehsenfeld	} NOAA Environmental Research Laboratories	Reactions of Charged Species
Dr. E. F. Ferguson		
Dr. D. R. Herschbach	} Harvard University	Homogeneous Chemistry, Kinetics and Spectroscopy
Dr. W. Klemperer		
Dr. L. E. Snyder	University of Illinois	Millimeter Wavelength Spectroscopy

December 31, 19

Table 5A
Research Funded by the Fluorocarbon Industry
and
Administered by the Manufacturing Chemists Association
WORK COMPLETED

Program	Investigator	Organization	Proposal Number	Completion Date
Measurement of fluorocarbons in the atmosphere*	Lovelock	U. of Reading	73-1	10/27/74
Monitoring of fluorocarbons in the atmosphere and simulation of atmospheric reactions of fluorocarbons	Taylor	U. of Calif.-Riverside	73-3	10/16/74
Investigation of spectroscopy of and photochemical changes in fluorocarbons	Sandorfy	U. of Montreal	73-2	10/11/74
Continuation of 73-1*	Lovelock	U. of Reading	74-3	12/31/75
Laboratory determination of sensitivity of laser-induced fluorescence for the detection of ClO under atmospheric conditions	Davis	U. of Maryland	74-10	5/31/75
Laboratory investigation of the feasibility of measuring ClO in the atmosphere by the chemical conversion-resonance fluorescence detection method	Stedman	U. of Michigan	74-7	2/28/75
Continuation of 73-3	Pitts	U. of Calif.-Riverside	74-2	12/31/75
Measurement of fluorocarbons and related chlorocarbons in the stratosphere and upper troposphere*	Rasmussen	Washington State U.	75-2	4/15/76
Laboratory and theoretical studies of the ultraviolet and visible electronic spectra of ClO*	Nicholls	York U.	75-11	6/14/76
Continuation of 74-2	Pitts	U. of Calif.-Riverside	75-12	4/15/76
Ground-based millimeter wavelength observations of stratospheric ClO	Ekstrom	Battelle Northwest	75-27	5/24/76
Investigation of the destruction of chlorofluoromethanes by naturally occurring ions	Campbell	Washington State U.	75-53	4/23/76
Development of an instrument to measure O, ClO, O ₃ , and total Cl in the stratosphere	Young	Xonics, Inc.	75-50	4/7/76
Investigation of ion-molecule reactions involving chlorofluorocarbons	Mohnen	State U. of N.Y.-Albany	75-64	4/1/76
Modeling of the fluorocarbon-ozone system*	Sze	ERT, Inc.	75-32	8/18/76
Critique of models used to estimate chlorofluorocarbon effects on ozone*	Cunnoid, Alyea, Prinn	CAP Associates	75-24	3/10/76
Studies of reactions of HO ₂ by laser magnetic resonance*	Thrush	U. of Cambridge	75-58	11/8/76
Continuation of 75-50	Young	Xonics, Inc.	75-86	11/15/76

*Work is continuing in a follow-on contract.

December 31, 1976

Table 5B
Research Funded by the Fluorocarbon Industry
and
Administered by the Manufacturing Chemists Association
WORK IN PROGRESS

Program	Investigator	Organization	Proposal Number	Contract Date	Contract Period
Measurement of reaction rates relevant to the fluorocarbon-ozone problem	Birks	U. of Illinois	75-1	4/17/75	15 mo. ^a
Measurement of stratospheric distribution of fluorocarbons and related species by infrared absorption spectroscopy	Murcay	U. of Denver	75-13	6/16/75	12 mo. ^a
Laboratory studies of the infrared vibration-rotation spectrum of ClO	Nicholls	York U.	75-30b	8/12/75	12 mo. ^a
Laboratory determination of the feasibility of laser magnetic resonance for ClO detection and reaction studies	Howard	NOAA-Boulder	75-47	11/13/75	12 mo. ^a
Collection and analysis of Antarctic ice cores	Rasmussen	Rasmussen Associates	75-84	11/29/75	3 mo. ^a
Measurement of fluorocarbon content of "antique" air samples	Rasmussen	Washington State U.	75-71	12/3/75	5 mo. ^a
Exploration for unidentified factors in the fluorocarbon-ozone problem	Lovelock	Private	75-67	12/4/75	12 mo. ^a
Laboratory measurement of spectroscopic absorption cross sections of ClO	Davis	U. of Maryland	75-73	2/5/76	8 mo. ^a
Continuation of program for ground level monitoring of ultraviolet solar radiation	Berger	Temple U.	75-62	2/5/76	12 mo.
Continuation of 75-2	Rasmussen	Washington State U.	75-59	2/11/76	9 mo. ^a
Measurement of OH in the stratosphere by laser induced fluorescence	Davis	U. of Maryland	75-87	2/24/76	12 mo.
Laboratory measurement of high resolution infrared spectra of chlorine-containing molecules of stratospheric interest	Murcay	U. of Denver	75-92	3/4/76	12 mo.
Laboratory investigation of the heterogeneous interaction of Cl and ClO with H ₂ SO ₄	Martin	Aerospace Corporation	75-81	3/22/76	12 mo.
Measurement of HCl, HF, ClO, etc., in the stratosphere by high resolution infrared spectroscopy	Girard	ONERA-France	75-88	3/29/76	12 mo.
Measurement of fluorocarbons and related chlorocarbons in the stratosphere by collection and analysis	Ridley	York U.	76-102	3/31/76	12 mo. ^b
Continuation of 75-13	Murcay	U. of Denver	76-101	4/1/76	12 mo. ^b
Construction of Fourier-transform spectrometer	Buijs	Bomem, Inc.	75-90	4/1/76	12 mo. ^b
Measurement of HCl and HF in the stratosphere by Fourier transform spectroscopy	Buijs	Bomem, Inc.	75-98	4/9/76	5 mo. ^a
Continuation of 75-32	Sze	ERT, Inc.	76-115	6/18/76	11 mo.
Total world ozone level: statistical analysis	Parzen, Pagano	Frontier Science and Technology Research Foundation, Inc.	76-106	7/15/76	12 mo.
The Electron Capture Detector as a Reference Standard in the Analysis of Atmospheric Halocarbons	Lovelock	Private	76-120	8/16/76	12 mo.
Continuation of 75-11	Nicholls	York U.	75-11 II	8/18/76	12 mo.
Continuation of 75-1	Birks	U. of Illinois	76-117A	8/26/76	12 mo.
Studies of Heterogeneous Reactions	Birks	U. of Illinois	76-117B	9/29/76	12 mo.
Studies of Compounds of Sulfur, Oxygen, and Chlorine	Kaufman	Emory U.	76-126	10/7/76	12 mo.

Table 5B (continued)

WORK IN PROGRESS					
Program	Investigator	Organization	Proposal Number	Contract Date	Contract Period
Microbiological and Multi-Dimensional Modeling Considerations Relating to Atmospheric Effects of Halocarbons	Cunnold, Alyea, Prinn	CAP Associates	76-122	10/8/76	12 mo.
Absolute Calibration of Fluorocarbon Measurements	Stedman	U. of Michigan	76-132	10/21/76	2 mo. ^a
Photochemical and chemical kinetic measurements of stratospheric importance with respect to the fluorocarbon issue	Timmons	Catholic U.	76-129	10/29/76	12 mo.
Atmospheric Chlorine Measurements in the Troposphere and Stratosphere	Eggleton	AERE Harwell	76-116	11/2/76	4 mo.
Laser Magnetic Resonance Study of NO ₂ Chemistry	Howard	NOAA-Boulder	76-100	11/6/76	6 mo.
Photochemistry of small chlorinated molecules	Wiesenfeld	Cornell	76-128	11/8/76	12 mo.
Atmospheric measurement of ClO and OH	Murcray	U. of Denver	76-135	12/9/76	4 mo.
Photochemical effects of fluorocarbons	Cunnold, Alyea, Prinn	CAP Associates	76-122S	12/10/76	12 mo.
Reactions of the NO ₂ radical studied by laser magnetic resonance	Thrush	U. of Cambridge	75-58 II	Pending	12 mo.
Far infrared wave observations of chlorofluoro- methane byproducts in the stratosphere	Solomon	SUNY Stony Brook	76-130	Pending	12 mo.
Far infrared stratospheric measurement of non-methane hydrocarbons	Rasmussen	Private	76-140	Pending	4 mo.
Far infrared laboratory comparisons of fluorocarbon measurements	Rasmussen	Private	76-142	Pending	3 mo.

^aContract extended.^bEstimate.

December 31, 1976

Table 6

PUBLICATIONS RESULTING FROM WORK SUPPORTED BY FLUOROCARBON MANUFACTURERS

Alexander Grant & Company

1. Environmental Analysis of Fluorocarbons FC-11, FC-12, and FC-22, February 5, 1976.

Allied Chemical Corporation

1. Statistical Modeling of Total Ozone Measurements with an Example Using Data from Arosa, Switzerland, W. J. Hill and P. N. Sheldon, *Geophys. Res. Lett.*, 2 (12), 541-4 (1975).
2. Analyzing Worldwide Total Ozone for Trends, W. J. Hill, P. N. Sheldon, and J. J. Tiede, *Geophys. Res. Lett.*, accepted.

J. W. Birks, University of Illinois

1. Four-Center Reactions Involving Dichlorine Monoxide, J.W.B., B. Shoemaker, T. J. Lock, and D. M. Hinton, draft ms.
2. Studies of Reactions of Importance in the Stratosphere. I. Reaction of Nitric Oxide With Ozone, J.W.B., B. Shoemaker, T. J. Lock, and D. M. Hinton, draft ms.

M. J. Campbell, Washington State University

1. Halocarbon Decomposition by Natural Ionization, M.J.C., *Geophys. Res. Lett.*, 3 (11), 661-4 (1976).

D. M. Campbell, F. N. Alyea, R. G. Prinn, Massachusetts Institute of Technology

1. The Impact of Stratospheric Variability on Measurement Programs for Minor Constituents, R.G.P., F.N.A., and D.M.C., *Bull. Am. Meteorol. Soc.*, 57 (9), 886-94 (1976).

E. I. du Pont de Nemours & Company, Inc.

1. Atmospheric Stability of Fluoroalkanes - Implications for Ozone Depletion, R. L. McCarthy and J. P. Jenson, Symposium on Fluorine Chemistry, Kyoto, Japan, August 26, 1976.
2. Measurement of the Reaction Rate of CF_3Cl with Atmosphere-Like Ions, R. G. Hirsch, *Atmos. Environ.*, 10 (9), 703-5 (1976).
3. Laboratory Microwave Spectrum of ClONO_2 and Evidence for the Existence of ClONO , R. D. Suenram, D. R. Johnson, L. C. Glasgow, and P. Z. Meakin, *Geophys. Res. Lett.*, 3 (10), 611-14 (1976), 3 (12), 758 (1976).
4. The Fluorocarbon-Ozone Theory. I. Production and Release, World Production and Release of CCl_3F and CCl_2F_2 (Fluorocarbons 11 and 12) through 1975, R. L. McCarthy, F. A. Bower, and J. P. Jenson, draft ms.
5. The Fluorocarbon-Ozone Theory. II. Tropospheric Lifetimes, An Experimental Estimate of the Tropospheric Lifetime of CCl_3F , J. P. Jenson, P. Meakin, and L. C. Glasgow, draft ms.

J. F. Lovelock, University of Reading

1. Atmospheric Halocarbons and Stratospheric Ozone, J.F.L., *Nature*, 252, 292-4 (1974) (11/22/74).
2. Long-range Transport of Photochemical Ozone in Northwestern Europe, R. A. Cox, A. E. J. Eggleston, R. G. Derwent, J.F.L., and D. H. Pack, *Nature*, 255, 118-21 (1975) (5/8/75).
3. Natural Halocarbons in the Air and Sea, J.F.L., *Nature*, 256, 193-4 (1975) (7/17/75).
4. Photochemical Oxidation of Halocarbons in the Troposphere, R. A. Cox, R. G. Derwent, A. E. J. Eggleston, and J.F.L., *Atmos. Environ.*, 10 (4), 395-8 (1976).
5. Halocarbon Behavior from a Long Time Series, D. H. Pack, J.F.L., G. Cotton, and C. Curthoys, draft ms.
6. The Electron Capture Detector Theory and Practice, J.F.L., *J. Chromatogr.*, 99, 3-12 (1974).

D. G. Murcray, University of Denver

1. Simultaneous Stratospheric Measurements of Fluorocarbons and Odd Nitrogen Compounds, W. J. Williams, J. J. Kusters, A. Goldman, and D.G.M., draft ms.

D. G. Murcray, University of Denver (continued)

2. Statistical-Band-Model Analysis and Integrated Intensity for the $10.8 \mu\text{m}$ Band of CF_2Cl_2 , A. Goldman, F. S. Bonomo, and D.G.M., *Geophys. Res. Lett.*, 3 (6), 369-12 (1976).
3. Measurements of Stratospheric Fluorocarbon Distributions Using Infrared Techniques, W. J. Williams, J. J. Kusters, A. Goldman, and D.G.M., *Geophys. Res. Lett.*, 3 (7), 379-82 (1976).
4. Measurement of Stratospheric Mixing Ratio Altitude Profile of HCl Using Infrared Absorption Techniques, W. J. Williams, J. J. Kusters, A. Goldman, and D.G.M., *Geophys. Res. Lett.*, 3 (7), 383-5 (1976).
5. Statistical Band Model Analysis and Integrated Intensity for the $11.8 \mu\text{m}$ Band of CFCl_3 , A. Goldman, F. S. Bonomo, and D.G.M., *Appl. Opt.*, 15 (10), 2305-7 (1976).

R. W. Nicholls, York University

1. The Absorption Cross-Sections and f-Values for the $v^n = 0$ Progression of Bands and Associated Continuum for the ClO ($A^2\Pi_1 + X^2\Pi_1$) System, M. Mandelman and R.W.N., *J. Quant. Spectrosc. Radiat. Transfer*, accepted.

J. N. Pitts, Jr., and O. C. Taylor, University of California at Riverside

1. Fluorocarbons in the Los Angeles Basin, N. E. Hester, E. R. Stephens, and O.C.T., *J. Air Pollut. Control Assoc.*, 24 (6), 591-5 (1974).
2. Relative Rate Constants for the Reaction of $\text{O}(^1\text{D})$ Atoms with Fluorocarbons and N_2O , J.N.P., H. L. Sandoval, and R. Atkinson, *Chem. Phys. Lett.*, 29 (1), 31-4 (1974) (11/17/74).
3. Reactions of Electronically Excited $\text{O}(^1\text{D})$ Atoms with Fluorocarbons, H. L. Sandoval, R. Atkinson, and J.N.P., *J. Photochem.*, 1 (4), 325-7 (1974).
4. Tropospheric and Stratospheric Chemical Sinks for Commercial Fluorocarbons, J.N.P. and R. Atkinson, *Trans. Amer. Geophys. Union*, 55 (12), 1153 (1974).
5. Mechanisms of Photochemical Air Pollution, J.S.P. and R. J. Finlayson, *Angew. Chem., Int. Ed. Engl.*, 14 (1), 1-13 (1975).
6. Fluorocarbon Air Pollutants. II, N. E. Hester, E. R. Stephens, and O.C.T., *Atmos. Environ.*, 9 (8-7), 603-6 (1975).
7. Fluorocarbon Air Pollutants, Measurements in Lower Stratosphere, N. E. Hester, E. R. Stephens, and O.C.T., *Environ. Sci. Technol.*, 9 (9), 875-6 (1975).
8. The Photostability of Fluorocarbons, S. Japar, J.N.P., and A. M. Winer, draft ms.
9. Background and Vertical Atmospheric Measurements of Fluorocarbons 11 and Fluorocarbon-12 over Southern California, L. Zafonte, N. E. Hester, E. R. Stephens, and O.C.T., *Atmos. Environ.*, 9, 1007-9 (1975).
10. Rate Constants for the Reaction of OH Radicals with CF_2Cl_2 , CF_2Cl_2 , and H_2 Over the Temperature Range 297-340K, R. Atkinson, D. A. Hansen, and J.N.P., *J. Chem. Phys.*, 61 (1), 1763-6 (1975) (4/1/75).
11. Tropospheric and Stratospheric Sinks for Halocarbons: Photo-oxidation, $\text{O}(^1\text{D})$ Atom and OH Radical Reactions, R. Atkinson, G. M. Brewer, J.N.P., and H. L. Sandoval, *J. Geophys. Res.*, accepted.
12. Fluorocarbon Air Pollutants. III. Fluorocarbon Measurements in the Lower Stratosphere, N. E. Hester, E. R. Stephens, and O.C.T., draft ms.

R. A. Rasmussen, Washington State University

1. Halocarbon Measurements in the Alaskan Troposphere and Lower Stratosphere, E. Robinson, R.A.R., J. Krasnee, D. Pierotti, and M. Jakubovic, draft ms.
2. Detailed Halocarbon Measurements Across the Alaskan Troposphere, E. Robinson, R.A.R., J. Krasnee, D. Pierotti, and M. Jakubovic, *Geophys. Res. Lett.*, 3 (6), 323-6 (1976).

C. Sanderfy, University of Montreal

1. Vacuum Ultraviolet and Photoelectron Spectra of Fluorochloro Derivatives of Methane, J. Doucet, P. Sauvageau, and C.S., J. Chem. Phys., 58 (9), 3708-16 (1973) (5/1/73).
2. Vacuum Ultraviolet Absorption Spectra of Fluoromethanes, P. Sauvageau, R. Gilbert, P. P. Berlow, and C.S., J. Chem. Phys., 59 (2), 762-5 (1973) (7/15/73).
3. Vacuum Ultraviolet Absorption Spectra of Chlorofluoromethanes From 120 to 65 nm, R. Gilbert, P. Sauvageau, and C.S., J. Chem. Phys., 60 (12), 4820-4 (1974) (6/15/74).
4. Vacuum Ultraviolet and Photoelectron Spectra of Fluoroethanes, P. Sauvageau, J. Doucet, R. Gilbert, and C.S., J. Chem. Phys., 61 (1), 391-5 (1974) (7/1/74).
5. On the Hydrogen Bond Breaking Ability of Fluorocarbons Containing Higher Halogens, T. DiPaolo and C.S., Can. J. Chem., 52 (21), 3612-22 (1974).
6. Fluorocarbon Anaesthetics Break Hydrogen Bonds, T. DiPaolo and C.S., Nature, 252, 471 (1974) (12/6/74).
7. Photoelectron and Far-Ultraviolet Absorption Spectra of Chlorofluoro-Derivatives of Ethane, J. Doucet, P. Sauvageau, and C.S., J. Chem. Phys., 62 (2), 355-9 (1975) (1/15/75).
8. Photoelectron and Far-Ultraviolet Spectra of CF_3Br , CF_2BrCl , and CF_2Br_2 , J. Doucet, R. Gilbert, P. Sauvageau, and C.S., J. Chem. Phys., 62 (2), 366-9 (1975) (1/15/75).
9. Photoelectron and Vacuum Ultraviolet Spectra of a Series of Fluoroethers, A. H. Hardin and C.S., J. Fluorine Chem., 5 (5), 435-42 (1975).
10. Ultraviolet Absorption of Fluorocarbons, a Review, C.S., Atmos. Environ., 10 (5), 363-51 (1976).

D. H. Stedman, University of Michigan

1. Measurement Techniques for the Ozone Layer, D.H.S., Res./Dev., January, 1976, pp. 22-4, 26.

N. D. Sze, Environmental Research & Technology, Inc.

1. Measurements of Fluorocarbons 11 and 12 and Model Validation: An Assessment, N.D.Z. and M. F. Wu, draft ms.

December 31, 1976



United Nations
Environment
Programme



Distr.
LIMITED

UNEP/WG.7/14(ABS)
2 March 1977

Original: ENGLISH

UNEP Meeting of experts designated by
Governments, Intergovernmental and
Non-Governmental Organizations on
the Ozone Layer

Washington, D.C., 1-9 March 1977

THE EFFECT OF CHLOROFLUOROCARBONS
ON THE ATMOSPHERE OZONE

Summary

The research program initiated in 1972 by chlorofluorocarbon manufacturers representing almost the entire world production exclusive of the U.S.S.R. and Eastern Europe is reviewed. Over \$3 million have been committed to date and a budget of over \$2 million has been approved for calendar year 1977.

Presently sponsored projects are discussed in the framework of the six NAS Panel recommendations for high priority research that could be accomplished in about two years. Comparisons are made with the earlier NAS Panel recommendations of July 1975 and those of the Interdepartmental Committee for Atmospheric Sciences of May 1975. Two documents are available that present the industry sponsored program in greater detail.^{1,2}

The results of recent work by Hill et al.³ on time series analysis of ozone data to detect abnormal trends are discussed briefly. It is suggested that this approach be examined thoroughly to determine its potential for (1) testing theoretical (model) predictions of ozone depletion and (2) providing a reliable "early warning system" for ozone depletion.⁴ If abnormal ozone trends can be determined by this technique with substantially greater sensitivity than was previously believed possible, practical limits can be placed on the scope of biological effects research and the urgency for regulation of chlorofluorocarbons as a precautionary measure will be diminished.

The results of Sze and Wu⁵ on model validation are discussed as an example of comprehensive analysis of experimental data employing alternative hypotheses. Data previously used to substantiate models in which an infinite tropospheric lifetime for FC-11 and FC-12 was assumed are shown to be equally consistent with lifetimes in the 10-30 year range. If the latter is true, the present most probable predictions of stratospheric ozone depletion by FC-11 and FC-12 will be reduced significantly. Care is urged in future comparisons of experimental measurements with model predictions.

The just announced⁶ finding at the National Bureau of Standards that sand and quartz promote photolysis of chlorofluoromethanes is noted. This observation is consistent with an early calculation by Rowland and Molina⁷ that indicated photolysis of C-Cl bonds might occur at wavelengths as long as 5000Å which can penetrate all the way to the earth's surface. The implications of these NBS results regarding adsorption and photolysis as a sink for FC-11 and FC-12 in the troposphere must be assessed quantitatively.

The chlorofluorocarbon industry welcomes comments and suggestions on its research program.

References and Notes

1. Summary, "Research Program on Effects of Fluorocarbons on the Atmosphere," Sponsored by the Fluorocarbon Industry, Revision No. 5, December 31, 1976.
2. Review of Industry-Sponsored Research, "The Effect of Chlorofluorocarbons on the Concentration of Atmospheric Ozone," December 31, 1976.
3. "Analyzing Worldwide Total Ozone for Trends," W. J. Hill et al., Geophys. Res. Lett. 4, 21-4 (1977).
4. The analysis indicates that the nine station network chosen for study allows detection of a 0.26% per year change over a six year period. This would detect a 1.56% total abnormal change and allow maximum depletion to be limited to 2.3%. It is estimated that an 18-station network would increase sensitivity to approximately a 1% abnormal change and this might be improved further by a larger network and satellite data.
5. "Measurements of Fluorocarbons 11 and 12 and Model Validation: An Assessment," N.D. Sze and M.F. Wu, Atmos. Environ. 10, 1117-25 (1976).
6. Chemical Week, March 2, 1977, p. 29.
7. "Chlorofluoromethanes in the Environment," F.S. Rowland and M. J. Molina, AEC Report No. 1974-1, Contract No. AT(043)-34, PA 126, September 5, 1974, pp. 17, 19.



United Nations
Environment
Programme



Distr.
LIMITED

UNEP/WG.7/15
25 February 1977

Original: ENGLISH

UNEP Meeting of experts designated by
Governments, Intergovernmental and
Non-Governmental Organizations on
the Ozone Layer

Washington, D.C., 1-9 March 1977

SWEDISH RESEARCH RELATED TO
THE STRATOSPHERIC OZONE PROBLEMS

Presented by

The Government of Sweden

DEPARTMENT OF METEOROLOGY
UNIVERSITY OF STOCKHOLM (MISU)
Henning Rodhe

1977-02-25

Sweden

Swedish research related to the stratospheric ozone problems

At present there is no research in Sweden aiming directly at the stratospheric ozone problems. There are, however, a number of research activities dealing with related topics, such as those listed below.

1. Department of Meteorology, University of Stockholm
(Head Prof B Bolin)
 - Precipitation chemistry (L Granat, R Söderlund)
 - Atmospheric chemistry of NO_x (R Söderlund)
 - Assessment of regional and global cycles of sulphur and nitrogen (L Granat, H Rodhe, R Söderlund)
 - Regional and global transport models (B Bolin, I Isaksen, H Rodhe, in cooperation with the Department of Geophysics, University of Oslo, Norway)
 - Measurements of O₃ and NO_x in the mesosphere (G Witt)
2. Swedish University for Agriculture, Forestry and Veterinary Medicine, Department of Microbiology (Head Prof B Norén), Department of Plant Biology (Head Prof S L Janson), Department of Forest Soils
 - Research about nitrogen cycling in soil and vegetation (S Janson, B Norén)
 - Denitrification processes (B Norén, H Nömmik)
 - N₂O-formation (U Granhall, T Rosswall, B Svensson)
 - N₂O-absorption in soils (B Svensson)
3. University of Gothenburg, Department of Dermatology (Head Prof G Swanbeck)
 - + Chromosomal damage to man from ultraviolet radiation
4. Swedish Aerosol Association, Box 238, 24020 Furulund, Sweden (B Bengtson)
 - Follows up producers research and development outside Sweden



**United Nations
Environment
Programme**



Distr.
Limited
UNEP/WG.7/16
1st March 1977

Original: ENGLISH

UNEP Meeting of experts designated by
Governments, Intergovernmental and
Non-Governmental Organizations on
the Ozone Layer

Washington, D.C., 1-9 March 1977

DANISH RESEARCH ACTIVITIES

RELATED TO THE HALOCARBON-OZONE PROBLEM - 1977

**Presented by
The National Agency of
Environmental Protection
Denmark**

NATIONAL AGENCY OF ENVIRONMENTAL PROTECTION

DANISH RESEARCH ACTIVITIES - RELATED TO THE 'GLOBAL WARMING-OZONE PROBLEM' - 1977 page 1

February - 77
UNEP/MG.7/16

Research field	Research projects (active)	Future projects (planned or proposed)	Institution	Researcher
Cancer	Cancer epidemiology. Watching the trend of the occurrence of different types of skin cancer in the total Danish population	Research of the causes of skin cancer by analysis of data of incidences related to e.g. age, occupation, place of living, smoking habits etcetera	Danish Cancer Registry, Copenhagen	J. Clemmensen
Climate	Study of climatic changes by Greenland ice-core analyses	Continuing	Geophysical Isotope Laboratory, University of Copenhagen	W. Dansgaard
Global radiation	Stochastic modelling of global radiation		Laboratory of Technical Hygiene - Technical University of Denmark, Lyngby	S. Hansen & N.B. Johansen
Halocarbons	Determination of lifetimes of simple halocarbons in the troposphere, based on theoretical estimates of rate constants	Extension to investigation of unsaturated and fully halogenated hydrocarbons	University of Aarhus, Institute of Chemistry	A. Grove & A.B. Mikkelsen
Halocarbons	An investigation of the stratospheric and tropospheric concentrations of halogenated aliphatic hydrocarbons in Greenland in the summer period	Continuing	Research Establishment Niso, Aerosol Sciences Laboratory, Roskilde	H. Flyver H. Zeuthen Høldsm & E. Lund Thomsen
Meteorology	Turbulent diffusion entrainment across inversion barriers	Continuing	Research Establishment Niso Meteorology Section, Roskilde	H.F. Busch
Nitrous oxide	Nitrogen fixation in association with mosses in fens and springs	Continuing	Botanical Institute, University of Aarhus, Århus	Esbern Wærncke
Nitrous oxide	Rates of denitrification in lakes as function of nitrate concentration, oxygen and temperature	Continued investigation of streams	University of Copenhagen, Freshwater Biological Laboratory, Hillerød	J.H. Andersen

DANISH RESEARCH ACTIVITIES - RELATED TO THE HALOGENATED-ORGANIC PROBLEM - 1977 page 2

Nitrous oxide	Nitrate reduction in the soil. Generation of N_2O and N_2 in different types of soil as a function of soil depth, soil oxygen and temperature	Continuing	State laboratory for soil and crop research, Lyngby	A. Lind
Nitrous oxide	Methods for measurements of denitrification (application of N_2)	Continuing	Water Quality Institute, ATV, Hørsholm, Denmark. Danish Isotopes Center, ATV, Copenhagen	K. I. Dahl-Jensen & J. Pfeiffer Madsen
Nitrogen modelling of atmospheric physics and chemistry	Development of accurate multidimensional numerical methods for simulation of atmospheric dispersion including chemical reactions	a) 1-D and 2-D dispersion and chemical modelling on global scale b) 1-D and 2-D climate modelling coupled to a)	Danish Meteorological Institute, Air Pollution Section, Copenhagen	L. P. Prabh & R. Gersonitz
Ozone in the stratosphere	Rocket sounding of ozone in the stratosphere, from Greenland, 1976		NASA, U.S.A. (Ionosphere Laboratory, Danish Meteorological Institute)	T. Tørnqvist
Photochemistry		Primary photochemical and photophysical processes of halocarbons. Primary thermal processes of fluorohalocarbons under atmospheric conditions	Institute for Chemistry, University of Copenhagen. Institute for Chemistry, University of Ålborg	O. Buchardt C. Lohse
UV radiation	UV radiation measurements in the Copenhagen area (Danish Meteorological Institute)	Continuing	Pennsylvania Temple University, (Finsen Institutet, Copenhagen)	H. Brodthagen
UV radiation	UV radiation measurements in Århus (WHO project)	Continuing	Århus University	Atmospheric Environment Service, Canada
UV radiation	Measurements of UV to infrared radiation and its effect on plant growth and development	Continuing	Royal Veterinary & Agricultural University, Copenhagen	A. Madsen

prepared by A. Astrup Jensen
& L. P. Prabh



**United Nations
Environment
Programme**



Distr.
Limited
UNEP/WG.7/17
1st March 1977

Original: ENGLISH



UNEP Meeting of experts designated by
Governments, Intergovernmental and
Non-Governmental Organizations on
the Ozone Layer

Washington, D.C., 1-9 March 1977

SURVEY OF OZONE RESEARCH IN NORWAY

**Presented by
Dr. Eigil Hesstvedt, Professor of Meteorology
Institute of Geophysics
University of Oslo, Oslo, Norway**

SURVEY ON OZONE RESEARCH IN NORWAY

presented by

Dr. Eigil Hesstvedt, professor of meteorology

Institute of Geophysics

University of Oslo, Oslo, Norway

1. OZONE RESEARCH BEFORE 1975

Ozone research has traditions in Norway. Observations of total ozone were started in 1939, and scientists interested in the climatology of atmospheric ozone are well acquainted with the long and nearly continuous record from Tromsø, situated at 70° North. Over a long time span this station provided important information about high latitude ozone. Shorter series of observations have been made at other stations: Dombås (62° N, 1940-1946) and Oslo (60° N, 1946-1949). Observations have also been made at Spitsbergen (79° N, 1950-1966). We have three Dobson photospectrometers at our disposal, but unfortunately none of these have been in operation since 1974. However, the instruments are now repaired and will be recalibrated. As soon as this process is finished, the instruments will be put in operation, one in Tromsø, one at Spitsbergen, and one in Oslo. (For details about ground based ozone measurements, see Appendix).

In addition to the ground based observations, the vertical distribution of ozone was measured in Oslo during the period 1968-1971 by means of ozone sondes. Altogether about 130 sondes were released. These observations constituted a part of the European ozone sonde network. However, the observations were abandoned

because of difficulties which arised from interference with local mobil-telephony. Further observations will depend on whether or not new radiocommunication equipment can be obtained. The future of ozone soundings from Oslo is therefore a question of economic resources and of training of personel.

While ground based ozone observations have long traditions in Norway, theoretical work, in the sense we use this notation today, only covers a period of about ten years. Furthermore, only few people have been involved.

Photochemistry of the upper atmosphere was taken up as a research topic at the Institute of Geophysics, University of Oslo. The research has been directed by Professor Eigil Hesstvedt, and the size of the staff has varied from one to three. In addition, well qualified research work has been done by a number of students as a part of their Master's Degree.

In the beginning, the research was directed at chemical characteristics of the mesosphere and lower thermosphere. One of the more important results of this work was the introduction of vertical eddy transport in the photochemical models. This changed our picture of the vertical distribution of chemical components with long photochemical lifetimes. Today the importance of this effect is generally accepted, and all models of today include transport mechanisms in one, two or three dimensions. In the late 1960's, our research was centered around ozone in the lower stratosphere. After the importance of nitrogen oxides was demonstrated in 1971 indepen-
dently by Johnston and Crutzen, our research followed a pattern similar to what we find in so many other institutes: a general study

of the chemistry of stratospheric ozone in order to evaluate the possible risk of a future fleet of high-flying aircraft. A series of papers were published about one- and two-dimensional modelling of the lower stratosphere. As an example, the first time-dependent photochemical model of the ozone layer was presented at the IAMAP General Assembly in Melbourne 1974. Much of this work fitted well into the goal of the Climatic Impact Assessment Program and was in part sponsored by the U.S. Department of Transportation. Our two-dimensional models were based on the best available knowledge of gas kinetics and stratospheric transport, and were used, among other things, to evaluate the effect of NO_x -injections from specified fleets and main routings of high flying aircraft (SST's). Figure 1 is taken from the Final Report (1974) of the CIAP and shows a comparison of various assessments of the SST effect.

2. OZONE RESEARCH AFTER 1975

Although we still shared an active interest in stratospheric problems, materialized in two institute reports, No.20 and 21 (see Appendix), our research has more and more become directed towards tropospheric ozone photochemistry. Our interest in tropospheric problems was initially raised a time after the successful oil drilling in the North Sea, when norwegian authorities were planning the development of petrochemical industry. In addition to inconveniences connected with industry in general, petrochemical industry involves emission of hydrocarbons and nitrogen oxides, which are the necessary ingrediences in photochemical smog formation. The analysis of the air pollution problems which could

arise in the surroundings of such industry was superficial, and it was felt that a better understanding of the mechanism of photochemical smog formation was needed. Against this background we decided to formulate a photochemical model which would cover the "clean" as well as the polluted atmosphere. A first version of the model was completed in the fall 1974. Later the model has been extended to include several different hydrocarbons which we observe in urban industrial areas.

Photochemical smog models is by no means a new invention; such models have been used in the analysis and forecasting of smog events. However, as a result of the complex dynamical and chemical character of the problem,, earlier models were based on very simplified chemical reaction scheme. In particular, chemistry of background air was simplified to a much larger extent than what is generally done in "clean" air modelling. The hydrocarbon chemistry often was treated by grouping reactions of the same type ("lumped" reactions). In contrast to this, our models make use of a "complete" background chemistry, i.e. we consider all known reactions which are likely to affect all the chemical components in the model. Furthermore, the hydrocarbon reaction scheme is complete in the sense that we consider explicitly the branching of the reaction scheme from the initial reactions of each hydrocarbon, through the intermediate steps to the end products.

It will appear from what we have said above that we may apply our model not only to simulate the chemistry of heavily polluted air, but also to describe the chemical composition of background ("clean") air and intermediate degrees of dilution which

occur when polluted air is transported away from its source and mixes with background air from the sides or from above.

I shall here describe briefly a model simulation of an atmospheric situation which serves as an example of present problem areas: photochemical oxidants, among them ozone, are known to occur in urban and industrialized regions almost all over the world. During the last years it has also become evident that these oxidants may in fact occur in high concentrations also when polluted air is transported into the environment and spread over large areas. The lifetime of ozone in such situations is long (of the order of days). To study such situations in more detail, we made the following model experiment:

We consider an air mass, with a prehistory of moderate emission rates, which is exposed to a somewhat stronger emissions of hydrocarbons and nitrogen oxides ($\text{HC}/\text{NO}_x = 1.5$). After a period of six hours the emission is abolished, and the further chemical development in the air mass is determined by pure photochemistry in one case and combined with dilution through mixing with overlying "clean" air in another case. The dilution is quantified by an increase in the mixing height according to the equation $\Delta H = \sqrt{4Kt}$, corresponding to air transport over the sea. Our experiment shows that ozone continues to increase with a marked diurnal variation for about 24 hours after it left the source region. During this period NO_x is almost completely converted to HNO_3 while the hydrocarbons, with the exception of acetylene, are broken down. Later ozone increases slowly at a steady rate. The time constant for the ozone decrease will vary strongly from one situation to the next, depending on the emission of precur-

sors in the source region, vertical mixing and the rate of deposition on the underlying surface. However, 3 weeks may be regarded as indicative for the chemistry, while mixing with "clean" air will reduce this figure to 1 week or less. However, our experiment shows, and this is supported by measurements, that anthropogenic production of ozone in polluted areas will affect the ozone levels on a regional scale. This brings us to the point where our modelling becomes relevant for the problem we shall discuss during this conference: anthropogenic changes in ozone on a global scale. A rough estimate of the magnitude of the anthropogenic production of ozone on a regional scale can be made as follows: the averaged flux of NO_x over a large area (Central Europe) is taken to be $2.5 \times 10^{11} \text{ molecules} \cdot \text{cm}^{-2} \cdot \text{s}^{-1}$. If we assume that each NO_x molecule on the average produces 2 ozone molecules (this is a low estimate) and a lifetime of one week, we arrive at an ozone increase of $3 \times 10^{17} \text{ molecules} \cdot \text{cm}^{-2}$ or 3% of the total ozone column density. This would apply to an area of the size of the industrialized part of a continent. A similar estimate of the increase in ozone averaged over the latitude belt $30-60^\circ\text{N}$ results in 0.25%. In this case the anthropogenic production of ozone is comparable ($\sim 1/10$) of the flux from the stratosphere.

Although these figures are relatively small, they are not insignificant. We feel that tropospheric ozone is not only an interesting topic in itself, but constitutes a part of the total ozone problem and as such deserves much attention in the future.

As was mentioned at the beginning of this survey, our research has also been directed towards stratospheric ozone.

Two reports were published last fall, one on a new method to compute the solar fluxes in the atmosphere and one, as a consequence of this, on the computation of more realistic values for photodissociation rate constants. The results of these two papers may be summarized as follows: Through the atmosphere, solar radiation is affected by a) absorption by atmospheric gases, b) molecular scattering, c) particle scattering, and d) reflection from clouds and from the ground. At the wavelengths of interest to stratospheric problems, O_2 and O_3 are the important absorbing gases. The reduction of solar fluxes due to absorption alone is easily taken care of when the total amount of O_2 and O_3 along the rays' path is known. The calculation is not very time consuming, and this effect is therefore included in the photochemical models where frequent recomputation of the fluxes due to a varying ozone distribution is necessary. On the other hand, the effect of multiple molecular (Rayleigh) scattering is more complex in its nature and therefore not so easy to quantify. Sundararaman et al. have solved the problem in its complete form and found that the effect is considerable and cannot be ignored. However, the computation is too time consuming to be included in atmospheric models. In an institute report from 1974 (No.5) we proposed an approximate solution of the problem by simply assume that the solar beam, which is scattered in all directions, is affected in such a way that one half of the scattered light proceeds in the direction of the beam, the other half in the opposite direction. The method was worked out in more detail in a recent report (No.20, October 1976) and tested against the complete solution

by Sundararaman et al. (see Figure 2). In spite of the extreme simplification, our method appears to approximate the real effect of scattering in a most satisfactory way, and since it requires a reasonable computer time, it is now possible to include the effect of multiple molecular scattering in modelling work.

The solar fluxes establish the basis for computations of the photodissociation rate constants (J-values) of a number of gases we consider in our models. Hence, realistic J-values can only be obtained if our flux data are realistic. Since all computations of possible ozone reduction from high-flying aircraft or from increased release of fluorocarbons referred to in the report from the National Academy of Sciences (September 1976) have ignored the effect of multiple scattering, it is of interest to see how much the modified J-values affect the ozone depletion estimates. Two gases, NO_2 and ClONO_2 , are of particular interest and the effect of including molecular scattering in the calculation of their photodissociation rate constants is demonstrated in Figures 3 and 4. We see that, in the stratosphere, the increase in solar fluxes due to multiple scattering results in a noticeable rate increase. As an immediate result we derive that the stratospheric content of chlorine nitrate (ClONO_2) has been overestimated in previous model calculations. We have tested the magnitude of this error in a one-dimensional model and find (Sunde & Isaksen, Report No.21) that the results of previous calculations of the possible ozone reduction due to fluorocarbons are to be multiplied by a factor of about 1.4.

3. PLANS FOR FUTURE RESEARCH

As already mentioned ground based observations of ozone are planned to start again in a near future in Tromsø, Oslo, and Longyearbyen.

The theoretical modelling work at the Institute of Geophysics, University of Oslo, will continue.

The models we have developed are very general in their scope and may be used to study problems of different kinds. Reports of work on tropospheric ozone are in progress, and we have formulated further problems to be studied later. We are also preparing a two-dimensional model of the troposphere and stratosphere. Even this model may be adopted to various problem areas.

Our major concern is the financial situation. Funding for atmospheric ozone studies are small at present, and the terms for two of the three staff members expire in about a year's time. In view of the importance of this part of the global environmental problem I express the hope that groups working in this field in Norway will be given a fair chance to make use of its scientific research potential to contribute to the international co-operation on the ozone problem.

APPENDIX

OBSERVATIONS OF ATMOSPHERIC OZONE IN NORWAY

(Compiled by Dr. Søren H.H. Larsen)

TOTAL OZONE

Total ozone has been observed in Norway intermittently since 1939 at three stations: Tromsø (70°N), Dombås (62°N), and Oslo (60°N). Two Dobson spectrophotometers, "old type", No.8 and 14, were used. The earliest series of observations were published in *Geophysica Norvegica*: Tromsø (1939-49, Tönsberg), Dombås (1940-46, Langlo), and Oslo (1946-49). Both instruments were recalibrated and modernized in 1949/50. Instrument No.8 was brought to Longyearbyen, Spitsbergen (79°N), where observations started in 1950 (S. Larsen). Attempts were made to use the moon as light source for observations during the polar night. The results from 1950-51 are published in *Geophysica Norvegica*. Data for the period 1951-66 were sent to the ozone world data center. In 1967 the instrument was moved from Longyearbyen to Ny-Ålesund, Spitsbergen, where only sporadic series of observations were taken. The instrument collapsed in 1973. This instrument, No.8, belongs to the International Ozone Commission.

The observations in Tromsø (instrument No.14) were continued until 1974. Data for the period 1950-74 were sent to the ozone world data center. It was difficult to get competent operators to the instruments in Tromsø and at Spitsbergen, and since 1975 both instruments have been out of operation.

Norsk Polarinstituttt made a series of observations in Antarctica (instrument No.56). The instrument was damaged and was left over to the Department of Physics, University of Oslo, where it has been repaired and calibrated against the standard instrument in Oxford.

The three instruments (No.8, 14, and 56) are now in Oslo. No.8 and No.14 are operational. No.14 will be sent back to

Tromsø where observations will be continued. Instrument No.8 will be sent to Longyearbyen where assistants at the meteorological office there will make the observations. No.56 will remain in Oslo, and new observations are planned. We also plan to make No.14 a standard instrument after intercomparison with another standard instrument.

VERTICAL DISTRIBUTION OF OZONE

Vertical distribution of ozone has been derived from optical ground-based observations in Tromsø (Tönsberg, Langlo) and at Spitsbergen (Larsen) (published in *Geophysica Norvegica*, 1945, and in *Nature*, 1954).

During the period 1968-71 a series of ozone sondes were flown from Oslo (Sören Larsen) under a project which aimed at the development of a combination of the MAST-sonde and the Väisälä meteorological sonde, which is used by the met. service in Norway. The results were satisfactory, but we had to make use of a Väisälä radioactively sonde to transmit the ozone signals. These sondes are not used by the Norwegian meteorological service. Observational data were exchanged with the European ozone sonde network and formed a part of the GARP experiment.

Difficulties with the telecommunications because of interference with mobil-telephoni has led to a temporary stop in the program. Our radiocommunication equipment has to be replaced in order to avoid interference. Ozone soundings require a well trained personal and considerable economic resources.

INSTITUTE REPORTS 1974-1977
INSTITUTE OF GEOPHYSICS (PHOTOCHEMISTRY GROUP)
UNIVERSITY OF OSLO

- No.6 On NO_2 -absorption of solar radiation and its role in the heat budget of the lower atmosphere (Hesstvedt, Isaksen,, August 1974)
- No.8 Formation of oxidants and other secondary pollutants in air mixed with nitrogen oxides and ethylene (Hesstvedt, November 1974) (also in *Geophysica Norvegica* Vol.31, No.2)
- No.9 Chemistry of a mixture of C_2H_4 , NO_x , and air, studied in a simplified photochemical dispersion model (Hesstvedt, March 1975)
- No.11 On the effect of chlorine on ozone formation in air polluted with C_2H_4 and NO_x (Hesstvedt, June 1975)
- No.15 On the filtering effect of atmospheric ozone (Hesstvedt, February 1976)
- No.16 On the chemistry of mixtures of hydrocarbons and nitrogen oxides in air (Hesstvedt, Hov, Isaksen, April 1976) (also in *Geophysica Norvegica*, Vol.31, No.6)
- No.20 A simplified method to include molecular scattering and reflection in calculations of photon fluxes and photodissociation rates (Isaksen, Midtbø, Sunde, Crutzen, October 1976) (also in *Geophysica Norvegica*, Vol.31, No.5)
- No.21 Stratospheric chlorine nitrate: reduced effect on the catalytical destruction of ozone due to molecular scattering and reflection of photon fluxes in the atmosphere (Sunde, Isaksen, November 1976)
- No.24 Diurnal variations of ozone and other pollutants in an urban area (Hov, Isaksen, Hesstvedt, January 1977)




FIGURE 1. Calculated Reduction of Vertical Ozone in Terms of Uniform Global Rate of NO_x Addition at 20 km (66 kft)
Source: Chang and Johnston, 1974
(taken from CIAP, Final Report)




FIGURE 2. Comparison of complete and simplified computation of solar fluxes




FIGURE 3. Rate constant for photodissociation of NO_2 :
broken line: pure absorption, solid line:
absorption and multiple scattering




FIGURE 4. Rate constant for photodissociation of ClONO_2 :
broken line: pure absorption, solid line:
absorption and multiple scattering



United Nations
Environment
Programme



Distr.
LIMITED

UNEP/WG.7/17(ABS)
1 March 1977

Original: ENGLISH

UNEP Meeting of experts designated by
Governments, Intergovernmental and
Non-Governmental Organizations on
the Ozone Layer

Washington, D.C., 1-9 March 1977

SURVEY OF OZONE RESEARCH IN NORWAY - A Summary

Presented by
The Government of Norway



United Nations Environment Programme



Distr.
LIMITED

UNEP/WG.7/17(ABS)
1 March 1977

Original: ENGLISH

UNEP Meeting of experts designated by
Governments, Intergovernmental and
Non-Governmental Organizations on
the Ozone Layer

Washington, D.C., 1-9 March 1977

SURVEY ON OZONE RESEARCH IN NORWAY

Norwegian measurements of total ozone were started in 1939 in Tromsø at 70°N. The record was discontinued in 1974. In shorter periods, measurements have also been made in other places: Dombås (62°N), Oslo (60°N) and at Spitsbergen (79°N). Norway has at its disposal three Dobson instruments which are now being repaired and recalibrated. These instruments will in the near future be placed in Oslo, Tromsø, and at Spitsbergen. Theoretical modelling work has over the last couple of years been concentrated on two topics: Tropospheric ozone, mainly in the lowest layers, and on calculations of solar fluxes and rates of photodissociation in the stratosphere and troposphere. Our tropospheric ozone model cover polluted as well as clean air and was primarily intended for computation of ozone production in urban and industrial areas (photochemical smog). Its flexibility has made it possible to study the chemical development after the polluted air has left its source region. We find the lifetime of ozone in such situations to be 1 - 3 weeks, depending on the conditions. Ozone is therefore spread over large areas without being broken down. Anthropogenic production of ozone in urban areas will therefore contribute to a raise (of the order 0, 1 - 1%) in total ozone. The main result of our stratospheric research is a new simplified method to include multiple molecular scattering in models. The solar fluxes, as computed in most models is thereby reduced to an extent which has a considerable impact on the photodissociation rates. An important effect is that on chlorine nitrate: The effect of releases of fluorocarbons upon total ozone, as compared in earlier models, are to be multiplied by a factor of 1.4 when multiple scattering is considered. Our research will be continued along the same lines. We have also developed a 2-D model for use in future studies.



**United Nations
Environment
Programme**



**Distr.
LIMITED**

**UNEP/WG.7/18
1st March 1977**

Original: English



**UNEP Meeting of experts designated by
Governments, Intergovernmental and
Non-Governmental Organizations on
the Ozone Layer**

Washington, D.C., 1-9 March 1977

**STATEMENT OF THE FEDERAL REPUBLIC OF GERMANY DELEGATION
CONCERNING RESEARCH IN THE FIELD
OF THE STRATOSPHERIC OZONE LAYER**

Presented by

Dr. Ulrici

Federal Ministry of Science and Technology

Statement of the German Delegation
Concerning Research in the Field of
the Stratospheric Ozone Layer

Mr. Chairman,
distinguished delegates,

First of all, on behalf of the delegation of the Federal Republic of Germany, I would like to thank the United States Government for providing the opportunity of meeting here in Washington, D. C. to discuss scientific matters in the field of the stratospheric ozone layer. We also express our gratitude for the initiative taken by the UNEP and the support provided by it.

The Government of the Federal Republic of Germany takes a serious view of the danger of a modification of the ozone layer as a result of the effects of certain halocarbons and other air pollutants and is therefore envisaging intensified endeavours in order to be able to contribute towards finding solutions to the scientific issues which have emerged. At present we can but confirm the opinion of the United States National Academy of Science, insofar as a great deal of uncertainty still prevails in this field which therefore needs a great deal of additional research.

The bulk of research work in the Federal Republic of Germany is presently concerned with the investigation of the presence, behaviour and effects of halocarbons. On the government side, a research program

covering the years 1976 to 1979 is currently being executed with promotion funds amounting to DM 4.7 million. Research projects requiring a further DM 2 million are in preparation. Other research activities are being carried out by industry to the order of approximately DM 1 million annually. The projects concern the improvement of analytical measuring methods, the measuring of vertical and horizontal profiles, as well as the North-South distribution of halocarbons with the aid of aircraft probes or registering balloons, and also laboratory tests of the photochemical reactions taking place in the atmosphere. Although priority is given to halocarbons, nitric oxides and other trace gases in the atmosphere are also detected at the same time in order to be able to base scientific conclusions on the evaluation of as comprehensive a range of data as possible.

Participants in the program are the universities of Freiburg, Wuppertal, Bonn, Kiel, Bochum, the Max-Planck-Institutes in Lindau and Mainz, the Nuclear Research Establishment Jülich, the Battelle Institute and the German Weather and Meteorological Service.

At Bonn University a project is also being carried out concerning the measuring of short-wave solar radiation, which is greatly influenced by variations in the ozone layer. This technique, which uses actinometric photometry, is to be carried out close to the earth in order to be able to correlate in the course of longtime experiments a possible biological effect with fluctuations in ultra-violet radiation. Another project

being carried out at Heidelberg University aims to detect natural sources of halocarbons by examining datable samples of ground water. The findings are to be combined to provide a trend analysis. The Max-Planck-Institute in Mainz is preparing a study on the metabolization of halocarbons in the biosphere.

The halocarbons program was launched in the Federal Republic of Germany in 1976. A symposium is planned to be held in Mainz in mid-1977, during the course of which the program will be presented and first results will be put up for discussion.

For the time being we have attached the greatest importance to the elaboration of the necessary basis of measuring data concerning the actual presence and behaviour of halocarbons. In the later phase of the program, which is to commence in autumn this year, we plan to include studies of the possible effects on the climate caused by halocarbons and other air pollutants. In particular, climate models are to be developed in order to provide some clues to the effects of emissions.

Moreover, studies are envisaged of the biological consequences, which after all are what interest us. We consider that it is all the more necessary to clear up the scientific uncertainty in this field because there is still a great deal of speculation on this very subject.

It will only be possible to investigate these issues at considerable cost and with some effort within a reasonable period. We are interested

in an efficient international division of efforts. We are prepared to make available the results of our work and to use our best efforts within this international framework in order to close gaps in knowledge.

In our view, the United Nations Environmental Programme is the appropriate forum for coordinating the handling of these global environmental problems in international cooperation. The Federal Republic of Germany is prepared to cooperate in the coordination of the scientific endeavours undertaken by the various governments and non-governmental organizations.

